

The Description of the 2324 Ring-dial His composed of two loncentric Circles of Silver or Brops the outer is the Meridian of our Horizon; y Inner is ye Equinoctial. The latter, in order to or present the Equator, is made moved upon two pivots, whereby it is fastoned tog Meridian, so as to be capable to traverse A at right Angles; & when'll brought to that Tituation, I meets with two Supporters, which propil, and priver its going fur ther; when ul is brought back with test, it falls. into two hotches, cut in the Axis, on purposes for its kaception. If we war to keep that Equator of the Elevation proper for each Horison, the Meridu must be suspended by a Circle, of thing which is brought to the Latitude of the place on y Meridian for if the time es et our grithe of suspension run on the Mexidian divided into Begrees, of the virlance of 49 begrees from

the Equator, that aing is at our Zenith. Therefore from y king by bole, there will, remain but 41 Begrees, since the bole en sequently of Equator of this I machine will be then at 41 begrees of Elevation on y Horizon; &y point of the pole at 49. These your arches To gether ese haust the iso begrees of the Horison, & the distance of the bole es always like y distance from y mith to the Equator. The Ring, to facilitate all the displacings nanted for every Horizon, Sink La double for loning in a Groove, which reaches along on the two chaces of y meridian. the siece of Suspension runs thus, as pleasure, as Jar as, under the northern polisand like the northern or Southern Latitude, regulating & position of theigh bouring pole, Arenders the King-dial on universal Instruments. The two poles are represented by two prots for tones to y Meridian Picle, or to the two potches onto which y Equinochial-civile is lowered

These two poles, or pivob, representing the poles of the world, support a Bridge which play in them by its Extremate and Cropses deams brically yo Equator, trought to its place to det its part, for the Equator is of no use, when Tolded in its Bose, where it breames concentric to the Meridian. The areis is represented by a long am harrow Aperture, which riggles along the middle of the Bridge. The use mod of this Aperluneis, lo todge in Na Iman prece of metal, which moves, with a grow through, called Cursor, & which running backwards & forwards under y Sun, according to the various declinations where Narriver from one Day to another, is found escaclly between the planet, & a point opposite on the inner Edge of y Equinochial; whene A follows, that the Jun, of Cursor, & if point opposite in the Equator of The King dial, boing on the Same Lim

WILEYBOURN

DESCRIPTION and USE

OFA

Portable Instrument,

Vulgarly known by the Name of

GUNTER'S Quadrant.

By which is perform'd most Propositions in Astronomy; as, the Altitude, Azimuth, Right Ascension and Declination of the Sun, &c. also his Rising and Setting, and Amplitude; together with the Hour of the Day or Night, and other Conclusions exemplify'd at large.

To which is added,

The Use of NEPIAR's Bones in Multiplication, Division, and Extraction of Roots; also the Nocturnal, the Ring-Dyal, and GUNTER's Line, in many necessary and delightful Conclusions, fitted to the Understanding of all Capacities.

By WILLIAM LEYBOURN.

The Third Edition, with the Addition of the Use of Gunter's Quadrant in taking the Declination of a Plain by the Help of a Horizontal Dial; also, the Use of a two Foot Rule in the Mensuration of Timber, &c.

By CHARLES LEADBETTER.

LONDON:

Printed for J. WILCOX, in Little-Britain, 1731.

983 William Courses the Course of the Course

to take the strengt Vitalia

and boild we have distributed

e el la participa de la company de la compan

HISTORICAL MENIC L

Car State Contract Co



The USE of a

Portable Instrument

Known by the Name of

GUNTER'S Quadrant.

in der geregen bei bei der geregen bei beite bei beite b

SECT. I.

Of the Description thereof.

Description of this Instrument; for all Persons that are capable of the Use thereof know, 'tis but

the fourth Part of a Circle, and divided in the Limb thereof into 90 Degrees, by equal Divisions, and number'd 10, 20, 30, 40, &c. unto 90; each Degree being subdivided into 4 Parts, above the Limb. There is inserted the twelve Months of the Year, and distinguished by capital Letters, viz. f. or fanuary, A 2

F. for February, M. for March, &c. and each Month is subdivided into 5 Days, more or less, according as the Largness of the Quadrant will bear; and then above the Days of the Month you have, in the Body of the Quadrant, the Hours and Azimuth Lines, together with Part of the Ecliptick and Horizon; also a Quadrant, &c. as will be much more obvious and intelligible to the Reader, by Inspection of the Figure thereof hereunto annexed, then by many Words.

2. And, to make the Quadrant compleat, there must be added a Couple of brass Sights with small Holes therein, sixed to one of the Semidiameters of the Quadrant; also a Thread or Silk sasten'd to the Center, with a Bead and Plumet as is usual: And the making hereof is well known to the Workman, and shew'd at large by Mr. Gunter. What else is necessary shall be explain'd

in its Use.

sulve les chain laire and

Lattinguilla bill por

SECT. II.

Of the Circle of Months.

His Circle of Months is figured with the Letters J. F. M. A. M. &c. fignifying the Months January, February, March, April, May, &c. each Month being divided unequally, ac-cording to the Number of the Days that are contain'd therein, as before is intimated.

1. And now to find any Day of the Month throughout the Year: First, fearch out the Month, and, from the Beginning thereof, account the small Divisions from one Month to another, each of them for five Days, (unless your Quadrant be large, and the Month divided into every Day) and by this Means you eafily find the Place (or Division) of the Day of the Month desired, which will be much easier by Practice, than can be express'd by Words: And here note, that every 10th Day is distinguish'd by a longer Line then the rest; thus, if I lay the Thread (being fixed in the Center) to the 10th Day of January,

'twill cut in about 21 Degrees in the Limb of the Quadrant: Now in June and December, when the Sun has his greatest North and South Declination, the Divisions here run close, and so the Day of the Month cannot be fo exactly found, but guess'd at, and near enough for most Uses; but in all the other Months of the Year, there needs not the least Scruple in this Particular. the to indmit of or gallace that are contain'd theren, as belore is

2. Having the Day of the Month, to find out the Sun's meridian Altitude, or bow high he will be at Noon?

Let the Thread be laid upon the Division call'd (or representing) the Day of the Month, and then that very Degree, which the Thread cuts in the Limb of the Quadrant, is the Sun's meridian Altitude required; fo the Thread, laid to the 15th Day of May, will cut 59 gr. 30 m. in the Quadrant's Limb, which is the Sun's Height that Day at Noon; and if it be laid upon the 18th Day, the meridian Altitude will be found 4 rgf. 39 m. Suredy based od yal I

Hiwi's

3. The meridian Altitude being known, to find the Day of the Month, which is the converse of the former.

The Thread, being laid upon the meridian Altitude, doth also fall upon the Day of the Month: So that if you have any one of them, you have both. So, if the Altitude at Noon be 59 gr. 30 min. (as before) and the Thread being set thereunto, it falls upon the 15th of May, and the 9th of July, and which of thefe two is the true Day, may be distinguish'd by the Quarter of the Year, or another Day's Observation: For, if the Altitude prove greater, the Thread will fall on the 16th Day of May, and 8th Day of July; or, if it prove lesser, the Thread will fall on the 14th Day of May, and the 10th of July; and thus the Doubt is fully determined, and by this we discover also what Days of the Year are of equal Length, which must needs follow by Consequence, if the meridian Altitudes be equal; and so the 4th of September and 18th of March have the same meridian Altitude, and the same Length, which is to be understood -1 Just L in

in the rest of the Months, as they stand

opposite in their proper Circles.

4. And here 'tis to be observ'd, that this Quadrant is accommodated but to one Latitude, and therefore (in this Case) serves but for one Place, and fuch which lye parallel thereunto, viz. East or West from thence, (as, suppose from the metropolitan City of London) those that lye 60 or 70 Miles North from this Parallel, the Sun will be a Degree lower, than appears by the Quadrant at Noon; and the contrary, one Degree higher from the meridian Altitude, to such as live 60 or 70 Miles South from London, or fuch Places that lye parrallel to it; this is to be understood if you be regulated by the Day of the Month upon the Quadrant.

5. But if you observe the Sun's Altitude just at Noon (or a little before, and a little after Noon) by holding the Quadrant in both your Hands, turning your left Hand towards the Sun, so that the Beams thereof may pass through the Holes of the Sights, then, the Plummet hanging at Liberty, the Degrees cut in the Limb by the Thread, is the

Alti-

call'd GUNTER's Quadrant. 9

Altitude desired in any Place; and the same may be done by the Moon and Stars, if you look through the Sights at the Moon or Stars, and observe then what Degrees are cut by the Thread in the Limb: So upon the 14th Day of April, just at Noon, the Sun-Beams passing through both the Sights, the Thread will fall upon 51 gr. 35 min. the true meridian Altitude of the Sun in the Latitude of London, for which Meridian most of the Quadrants are made.

SECT. III.

Of the Hour-Lines on the Quadrant.

on the Center of the Quadrant to the Beginning of the Sun's Declination, doth represent the Equator, or equinoctial Circle (and is usually made a Radius or Tangent) of 45 Degrees; but that Arch which is drawn by 23 gr. 30 min. of Declination, and is just above the Circle of Months and Days, represent the Tropicks of Capricorn and Cancer

Cancer, or the Sun's greatest Declination both North and South; and those Arch-Lines, which are described upon the Quadrant between the Equator and the Tropicks, being undivided and number'd at the Equator by 6, 7, 8, 9, 10, 11, 12, and at the Tropicks by 1, 2, 3, 4, 5, 6, 7, 8, those very Lines or Arches are the Hour-Circles.

2. That which is drawn from 12 in the Equator to the Middle of June, represents the Hour of 12 in the Summer; and those which are drawn with it towards the right Hand, are the Hours of the Day in Summer, and also of the Night in Winter: But that Arch which is drawn from 12 in the Equator to the Middle of December, represents the Hour of 12 in the Winter; and those Arches with it to the left Hand, the rest of the Hours of the Day in Winter, and Night in Summer; and of both these, that which is drawn from 11 to 1, serves for 11 in the Forenoon and 1 in the Afternoon; that which is drawn from 10 to 2, ferves for 10 in the Forenoon and 2 in the Afternoon: The Reason is, that the same Day the Sun has the same Altitude 2 Hours before Noon, as it has 2 Hours af-

call'd Gunter's Quadrant. 11

ter Noon; and the same Reason is to be understood in all the rest, which is plain and easily understood, as by the sollowing Propositions will clearly appear.

3. The Day of the Month, or meridian Altitude of the Sun, being given, to find the Place of the Sun in the Ecliptick.

Let the Thread be laid to the Day of the Month, or to the Sun's Height at Noon, (for the one gives the other) obferve then where it crosses the Hour of 12, and fet the Bead to that Intersection; then move the Thread till the Bead fall on the Ecliptick Circle, (which is known by the Characters of the 12 Signs) and it will fall upon the Sun's Place; and fo if it were the 15th of May, or meridian Altitude 59 gr. 30 min. lay the Thread as aforesaid, and put the Bead to the Intersection of the Hour of 12, and then move the Thread 'till the Bead fall on the Ecliptick, and it will there touch upon the 4th Degree of Gemini, and the same of Sagittary; also the 26th of Cancer and the 26th of Capricorn; and which of these is the Place of the Sun, may appear

appear by the Quarter of the Year, or another Day's Observation.

4. To restify the Bead for the Hour of the Day, &c.

Knowing the Day of the Month, place the Thread thereon, and then, slipping the Bead to the Meridian or Hour of 12, for the Season of the Year, you are fitted to find the Hour by the Sun; but for the Stars, only lay the Bead upon the Star proposed. And thus, the Bead being rectify'd, if you do but obferve the Altitude of the Sun, the Bead will fall on the Hour-Line defired or fought, if you bring the Thread to the Altitude: So on the 10th of April, having rectify'd the Bead for the Time, you may find the Altitude 36 gr. at the same time the Bead falls upon the Hour-Line of 3 and 9; and then I may conclude, 'tis 9 in the Forenoon, or 3 in the Afternoon; and so upon the 9th of April, the Bead being rectify'd, and the Altitude obferv'd 3 gr. the Bead will fall between 8 and 9 in the Equator, and between 3 and 4 in the Tropicks; so that 'tis half an Hour past 8 in the Morning, or half an

call'd GUNTER's Quadrant. 13

Hour past 3 in the Afternoon, and here observe, if the Sun ascends, it is be Forenoon; but if the Sun descends or falls in Altitude, 'tis Afternoon, as Reason itfelf will direct: So again, upon the aforefaid 10th of April, if the Sun's Altitude be 40 gr. you will find the Bead at the fame time fall half way between the Hour-Lines 9 and 10, or 2 and 3; wherefore it must either be half an Hour past 9 in the Morning, or half an Hour past 2 in the Afternoon, which is most easily distinguish'd. And after the same Method, if upon the 1st of November I obferve as aforefaid, and find the Bead to fall half way between 8 and 9, and 3 and 4 in the Winter-Hour-Circles; hence I conclude, 'tis half an Hour past 8 in the Forenoon, or half an Hour past 3 in the Afternoon, & sic de cæteris.

5. The Place of the Sun in the Ecliptick being known, to find the Day of the Month.

Lay the Thread and Bead on the Place of the Sun in the Ecliptick, and then move the Thread and Bead to the Hour of 12, and so, if the Place of the Sun be 4 gr. of Gemini, the Bead being laid to this Degree, and then moved to the

Hour

Hour of 12 in Summer, the Thread will fall on the 15th Day of May, or the 8th of July; or, if it be moved to the Hour of 12 in Winter, the Thread will fall on the 6th of January and the 14th of November, and which of these is the Day of the Month required, the Quarter of the Year will determine.

By this and the third Proposition, there is two Ways shew'd to rectify the Bead by the Place of the Sun, and by the Day of the Month; but the best Way is that by the Place of the Sun, for in the other some small Difference may arise by the Leap-year. There is also a third Way by a Table of the Sun's Declination for each Day of the Year, (which Seamen are seldom without) and if so, the Bead may be set thereunto in the Line of Declination.

6. To know how high the Sun will be, at any Hour of the Day given, throughout the Year.

The Bead being rectify'd for the Time, by any of the Ways aforesaid, bring it to the Hour given from the Hour of 12 towards the Line of Decli-

nation

call'd Gunter's Quadrant. 15

nation, and the Degrees, cut by the Thread in the Quadrant or Limb thereof, will shew the Altitude of the Sun at that Time; so upon the 30th of March, at 8 in the Morning and 4 in the Afternoon, the Sun will be found to be 24 gr. 30 min. high: Again, on the 10th of April, the Sun being in the Beginning of Taurus, the Bead being rectify'd, you shall find the Altitude at Noon at London 50 gr. 15 min. but at 11 in the Morning 48 gr. 12 min. at 10 of the Clock but 43 gr. 12 min. at 9 only 36 gr. at 8 of the Clock 27 gr. 30 min. at 7 but 18 gr. 18 min. at 6 in the Morning but 9 gr. and at 5 a Clock it touches the Line of Declination, and has no Altitude at all; whence we may conclude, the Sun did rife much about that Hour. And then, if you move the Thread again from the Line of Declination towards the Hour of 12, you shall find the Sun 8 gr. 33 min. below the Horizon at 4 in the Morning, and near 16 gr. at 3, and 21gr. 51 min. at 2, and 25 gr. 40 min. at 1, and 27 gr. at Midnight. For the Bead being rectify'd as aforesaid, and brought to the Hour of the Night (which, as before was intimated, are the Winter-Hours B 2 in

in the Summer, and the Summer-Hours in Winter) then the Thread in the Limb cuts the Degree of the Sun's Depression for any Hour of the Night; and thus 'twill be found, that at a 11 of the Clock at Night, the 30th of March, the Sun will be found at least 28 gr. 30 min. below the Horizon.

7. The Hour of the Night given, to find the Sun's Depression.

And here let it be observed, that the Sun is always as much below the Horizon at any Hour of the Night, as his opposite Point is above the Horizon at the like Hour of the Day, and therefore (to make this more plain) the Bead being fet, if the Question be put for any Hour of the Night in Summer, then move it to the like Hour of the Day in the Winter-Hours: But if of any Hour of the Night in Winter, then move the Bead to the like Hour of the Day in Summer, and so the Degrees, cut by the Thread in the Limb of the Quadrant, shall shew how much the Sun is below the Horizon at that Time: And to suppose it the 10th of April, at 4 of the Clock in the

the Morning, the Bead being set to his Place, according to the Time in the Summer-Hours, and brought to 4 of the Clock in the Afternoon in the Winter-Hours, you will find the Thread to cut 8 gr. and about 30 min. in the Limb of the Quadrant, which is the true Depression of the Sun at that Time.

8. The Depression of the Sun supposed, to give the Hour of the Night with us, or the Hour of the Day to our Antipodes.

In regard the Sun is as much above the Horizon at all Hours of the Day, as his opposite Point is below the Horizon at the like Hours of the Night; therefore, first set the Bead according to the Time, and then bring the Thread to the Degree of the Sun's Depression below the Horizon, fo shall the Bead fall on the contrary Hour-Lines, and thereby shew the Hour of the Night in respect of us, which is the like Hour of the Day to our Antipodes: So the 10th of April, the Sun being then in the Beginning of Taurus, and, by Supposition, 8 gr. 30 min. below the Horizon in the East, it is required to be known, What Time of B 3

the Night it is? First set the Bead according to the Day in the Summer-Hours, then bring the Thread to 8 gr. 30 min. in the Limb of the Quadrant, so shall the Bead fall among the Winter-Hours on the Line of 4 of the Clock in the Afternoon; wherefore to our Antipodes 'tis 4 of the Clock in the Afternoon, and to us 4 in the Morning.

9. Having the Time of the Year, or the Place of the Sun being known, how by this Quadrant to find the Beginning of Day break, and End of Twilight.

There is very little Difference between this Proposition and the former; for the Day is said to break when the Sun comes to be but 18 gr. below the Horizon in the East, and Twilight to end when the Sun is 18 gr. below the Horizon in the West; wherefore let the Bead be set for the Time, and then bring the Thread to 18 gr. in the Quadrant, so shall the Bead, falling upon the contrary Hour I ines, there shew the Hour of Twilight, as before in the Operation of the last Proposition, and so 'twill be found a little more then a Quarter after

call'd Gunter's Quadrant. 19

3 in the Morning, for Day-break upon the 10th of April; after the same Manner upon the 30th of March, Day-break will be found to be at 3 of the Clock in the Morning; and if so, we may easily judge when 'twill be dark at Night, which is said to be as much after 6, as this is before 6, and then we may conclude 'twill be dark at 9 of the Clock at Night, it being then past Twilight.

SECT. IV.

Of the Azimuth-Lines.

which are drawn between the Equator and the Tropicks, on that Side of the Quadrant towards the right Hand, or nearest to the Sights, and are number'd with 10, 20, 30, 40, &c, in the Equinoctial, then downwards from the Equator to the Tropicks with 110, 120, 130, &c. the uttermost towards the left Hand representeth the Meridian; that which is number'd with 10, the tenth Azimuth from the Meridian;

dian; and that which is number'd with 20, the twentieth; and so of the rest. Those Lines, which are drawn from the Equator to the left Hand, do shew the Azimuth in the Summer; and those other to the right Hand, do shew the same in the Winter. The Uses of them follow.

2. The Azimuth, the Sun is upon being known, to find the Altitude of the Sun above the Horizon.

First let the Bead be set for the Time, as before is shewed, then move the Thread until the Bead fall on the Azimuth; fo the Degrees, which the Thread cutteth in the Quadrant, shall shew the Altitude of the Sun at that Time: Where: you are to observe, that, in regard the Azimuths are drawn on the right Side: of the Quadrant, you are also to begin, to number the Degrees of the Sun's Altitude, from the right Hand towards the left, contrary to the Directions of some: former Propositions. Examples will clear this Point: Suppose the Time given were the fecond of August, when the: Sun hath but 14 gr. 49 m. of North Declination,

call'd Gunter's Quadrant. 21

clination, set the Bead for the Time, and the Sun's Meridian Altitude, or Height at Noon, will be found 53 gr. 17 min. But when he is 10 gr. from the South, 53 gr. 10 min. when 20 gr. then about 52 gr. 8. min. when 30 gr. then 50 gr. 20 min. when 40 gr. then 47 gr. 28 min. when 50 gr. then 44 gr. 12 min. when 60 gr. then 39 gr. 35 min. when 70 gr. then 33 gr. 50 min. when 80 gr. then 27 gf. when the Sun is in the East or West 90 gr. from the Meridian, then is the Altitude near 19 gr. 20 min. when 110 gr. then 3 gr. 20 min. and before he cometh to the Azimuth 120 gr. he has no Altitude, for the Sun having 14gr. 49 min. of North Declination will rise and fet at 114 gr. 34 min. Azimuth from the Meridian.

3. The Altitude of the Sun being given, to find on what Azimuth, or Point of the Compass, he bears from us.

Set the Bead in his proper Place for the Day proposed (as before directed) and then, observing the Sun's Altitude, lay the Thread to the Complement of that Altitude, (which is but accounting

so!

so many Degrees from 90 gr.) on the Quadrant, so shall the Bead fall upon the true Azimuth required. Example: Suppose upon the second of August (having rectify'd the Bead) you observe and find the Sun's Altitude to be 19 gr. 20 min. then remove the Thread to the Complement thereof, viz. 70 gr. 40 min. or, which is all one, upon 19 gr. 20 min. accounted from 90 gr. from the right Hand to the left, and the Bead will fall upon 90 gr. from the Meridian; and therefore the Point the Sun is upon bears from us, either full East or full West; and which of these it is, may immediate. ly be discovered by a second Observation; for if the Sun, increase in Altitude, 'tis Forenoon, and the Azimuth East; if the Sun decrease in Altitude, 'tis Afternoon, and then the Azimuth is West; which is as easily discovered, by the Hour of the Day. And so again, if I find the Sun's Altitude to be 25 gr. if I lay the Thread to 25 gr. accounted back-ward from 90 gr. the Bead will fall upon very near 80 gr. from the South, and that is the Point of the Compass the Sun is upon Eastward, if in the Morning; and Westward, if in the Afternoon. By these

call'd Gunter's Quadrant. 23

these Propositions you may know the Altitude of the Sun every Hour of the Day, though he appears not; and what Part of the Heavens he is in likewise; and, then knowing what Azimuth the Sun beareth from us, you may find not only a Meridian Line, but the Coasting of a Country, the Situation of a Building, and the Variation of the Compass, &c. as for Example: Suppose some Day in the Afternoon, I should find, by the aforesaid Direction, that the Sun bears from me 60 gr. from the South towards the West, then, there being 90 gr. in each Quarter of the Compass, the West will appear to be 30gr. to the right Hand, the East is just opposite to the West, and the Meridian-Line, or North and South, lyes in the midst between them.

4. To find the 4 Cardinal Points upon an horizontal Plain.

First find the Azimuth of the Sun, and instantly lay your Quadrant upon the Plain, so that the Face thereof may lye uppermost, then hold in your Hand some small String with a Plummet hanging thereunto, and move the Quadrant till

the Sun casteth the Shadow of the String through the Degree of the Limb which was found for the Azimuth, and through the Center of the Quadrant; then, holding the Quadrant still, draw Right-Lines by each Side thereof, and protract them, so that they may cross each other, and these Lines shall point out East, West, North, and South.

5. If two several Sun-Shadows be observed, the one in the Forenoon, and the
other in the Afternoon of the same Day,
exactly to touch with their Ends, the
Circumference of the same Circle describ'd
in a Plain, parallel to the Plain of the
Horizon, the Line from the Center, equally dividing the Arch of that Circle betwist the two Shadows, will be the true
Meridian in that Place.

This is a useful Theorem, and may be better understood by Example, viz. Take a Piece of Board or Mettle, and let it be so evenly placed, that it may lye parallel with the Plain of the Horizon; in this Plain describe divers Circles from the same Center, and in that Center raise a Gnomen at Right-Angles.

This

call'd Gunter's Quadrant. 25

This Platform being thus order'd, let the Shadow of the Gnomen, or Wire, be observ'd some time before Noon, till such time it do exactly touch the Circumference of one of these Circles. Again, in the Afternoon, let the Shadow be observed till the End of this Perpendicular touch the Circumference of the same Circle, which will exactly happen so many Hours after Noon, as the other before Noon; then you are diligently to observe these two Points in the Arch of the faid Circle, and divide that Arch into two equal Parts by a Line from the Center; this very Line is a true meridian Line, upon which when the Shadow falls, you may be fure 'tis just 12 of the Clock, or just high Noon.

6. To find the Declination of a Wall for the making of a declining Dial.

Apply one Edge of the Quadrant to the Wall, and at the Limb hold up a Thread and Plummet, so that the Shadow thereof may pass through the Center, then, marking what Degree of the Limb the Shadow of the String so held

shall touch, presently find the Degree in the Limb of the Sun's Azimuth at that time; then, if you have done right, the Degrees between the Degree of the Azimuth of the Sun, and the Degree of the Shadow of the String so mark'd, shall be the Degree of the Declination of the Wall from the true Meridian; which, whether it be Eastward or Westward, may be known by comparing it with the Sun. And note, this Proposition may be the better perform'd, the farther the Sun is from the Meridian.

7: To find how much a Plain or Wall declines from the Zenith.

Apply the Edge of the Quadrant to the Plain or Wall, so that the Center may be downwards, and the Limb upwards; then hold a String and Plummet, so that the Thread or String may touch the Center and the Limb, and number the Degrees between the Plain and the String, which will shew the Quantity in Degrees of the Plain's Reclination

call'd GUNTER's Quadrant. 27

8. To find the Elevation of the Pole in any Place.

First take the meridian Altitude of the Sun by Observation, (or otherwise) and also the Sun's Declination (as shall be shew'd) which if it be Northwards, substract, if Southwards, add to the Sun's meridian Altitude, and you have the Complement of the Pole's Elevation, or the equinoctial Height; which subducted out of 90 gr. leaves the Latitude of the Place desir'd, which is so plain, it needs no Example.

SECT. V.

Of the Ecliptick Circle or Arch.

guished from the other Circles, and is represented by the Arch, to which is annexed the Characters of the 12 Signs, and is drawn from the Hour-Line of 6 in the Equator, to the other side of the Quadrant, ending in the Tropicks of

Cancer and Capricorn, each Sign being divided unequally into 30 gr. which is to be reckon'd both ways from the Characters of the 12 Signs, Aries, Taurus, Gemini, Cancer Leo, Virgo, Libra, Scorpio, Sagittary, Gapricorn, Aquary, Pisces, which Signs do also answer to the 12 Months; and the Thread being laid upon the Day of the Month, and the Bead rectify'd, or brought to the proper Meridian, or 12, as before has been exemplify'd, according to the Season of the Year, viz. to the Summer-12 in Summer, and the Winter 12 in Winter, &c. Then bring the Bead to the Ecliptick (fo rectify'd) and it will fall upon the Sun's Place therein for that Day, regard being had to the Time of the Year: So upon the 18th of March, the Sun's Place, by the falling of the Bead, will be about 8 gr. of Aries, and upon the

17th of October in 4 gr. 30 m. of Scorpio.

2. In finding the Place of the Sun in the Ecliptick, 'tis necessary to know which Way the Sun declines from the Equator; for if he declines North, the Character belonging to his Place is above the Ecliptick, if South, beneath it. Now, for as much as the Bead will

fall

call'd GUNTER's Quadrant. 29 fall between two Characters, you may know which it is, by a fecond Day's Observation; as for Example: Let the 20th of August be suppos'd, the Thread being plac'd upon the Day, bring the Bead to the Summer-Meridian, which will shew that the Sun is then in its Northern Declination (because the Bead very plainly shews the same) and the Thread being moved falls either on the 7th gr. of Virgo, or at 22 gr. 3 m. of Aries, which the Time of the Year will plainly distinguish; for Aries, Taurus, Gemini, Cancer, Leo, Virgo, are the Northern or Summer-Signs, and Libra, Scorpio, Sagittary, Capricorn, Aquary and Pisces, are the Southern or Winter Signs: And this Proposition will be found ex-

ceeding easy by a little Practice.

3. If the Thread be laid upon the Degree of the Sun's Place in the Ecliptick exactly, it will cut in the Limb of the Quadrant, the Right Ascension of the Sun; so the Sun's Place being 4 gr. of Gemini, the Thread laid upon this Degree, will cut 62 gr. in the Limb for the Sun's Right Ascension requir'd: But if the Sun has pass'd about 90 gr. from Aries, or more than 3 Signs, then there must

be more than 90 gr. allow'd to the Right Ascension: Thus, if the Degrees of the Sun's Place be between Aries and Cancer, then his Right Ascension is the very Degrees cut by the Thread in the Limb; but if the Sun's Place be between Cancer and Libra, then to the first Quadrant you must add those Degrees cut by the Thread in the Limb, counting backwards from the Sights-side; if between Libra and Capricorn, then to 180 gr. add what the Thread cuts; if between Capricorn and Aries, then to 270 gr. add the Degrees cut by the Thread in the Limb, counting backwards again, and you have the Sun's true Right Ascension defired: So if the Sun be in 26 gr. of Cancer, though the Thread falls upon 62 gr. as before, yet the Complement thereunto must be added, viz. 28 gr. and so the Right Ascension will be 118 gr. So upon the 18th of March, the Right Afcension is but 7 gr. but upon the 5th of August 145 gr. upon the 10th of October 206 gr. and upon the 9th of March 360 gr. and thus the Business of Right Ascension is made as plain as possible may be to the meanest Capacity. Usus promptus facit.

And

call'd GUNTER's Quadrant. 31

And thus, having the Right Ascension given, you may by the converse Work find the Sun's Place; for lay but the Thread to the Sun's true Right Ascension in the Limb, and it will cross the Sun's Place in the Ecliptick desired.

Markararararar arabarararararar

SECT. VI.

Of the Line of the Sun's Declination.

HE Line of Declination is drawn from the Center to the Beginning of the Quadrant, and divided from the Beginning of Aries downwards to 23 gr.

29 min.

Its Use is thus, rectify the Bead as before shewed, and bring it to the Line of Declination, and you have the Sun's Declination for that Day: So if the Sun's Place given be the 4th gr. of Gemini, the Bead sirst set to this Place, and then moved to the Line of Declination, will there shew it to be 20 gr. 59 min. Declination, Northerly from the Equinoctial; and so, if you have only the Day of the Month given, you may presently

32 The Use of a portable Instrument, sently find both the Sun's Place, and Declination also for that Day: Thus upon the 17th of October the Sun has 13 gr. of Declination South. And note, that if

the 17th of October the Sun has 13 gr. of Declination South. And note, that if the Bead fall upon the Winter-12, the Declination is South; but if upon the Summer-12, 'tis North; and, having the Declination only given, you may thereby find the Sun's Place, and Day of the Month.

MANAMANANANANANANANANANANANANANANAN

S E C T. VII.

Of the Horizon.

upon the Quadrant by the Arch drawn from the Beginning of Declination towards the End of Frebruary, divided unequally, and number'd by 10, 20, 30, 40, &c.

2. Having the Place of the Sun, or Day of the Month, to find the Amplitude of the Sun's Rising and Setting.

call'd Gunter's Quadrant. 33

Let the Bead, rectify'd to the Time given, be brought to the Horizon, and there it shall shew the Amplitude required: So if the Day of the Month given be the 15th Day of May, and the Sun the 4th gr. of Gemini, the Bead, rectify'd and brought to the Horizon, shall there fall upon 35 gr. 8 min. and this is the Amplitude of the Sun's Rising from the East, and Setting from the West; which Amplitude is always North if the Sun be in northern Signs, but when he is in fouthern Signs, then 'tis always Southward; that is, from the 10th of March to the 11th or 12th of September North, but all the other half Year, the Amplitude of the Sun is Southerly; and so upon the 9th of April, the Amplitude will be found just 18 gr. Northerly. Understand the same Method in other Operations.

3. The Day of the Month, or the Sun's Place given, to find the Sun's Ascensional Difference.

Let the Bead, rectify'd for the Time, be brought to the Horizon, so shall

34 The Use of a portable Instrument,

the Degrees, cut by the Thread in the Quadrant, shew the Sun's Ascensional Difference: So upon the 10th of May, the Sun being in the last Degree of Tau-rus, or Beginning of Gemini, let the Bead be rectify'd and brought to the Horizon, and then the Thread, in the Limb of the Quadrant, shall shew the Ascensional Difference to be 27 gr. 30 min. And so upon the 15th of May, the Sun being about 4 or 5 gr. of Gemini, the Ascensional Difference will be found near 29 gr. or rather 28 gr. 50 min. So upon the 9th of April the Ascensional Difference will be found 15 gr. Now, having this, you may thereby find the Time of the Rising and Setting of the Sun, (which by some of the former Propositions may be guess'dat) but more accurately thus; if the Ascensional Difference be converted into Time, by allowing one Hour for every 15 gr. and 4 min. of an Hour for each Degree, and this will shew how long the Sun riseth before six in the Summer (that is, from the 10th of March to the 11th or 12th of September) and after six of the Clock in Winter: So if the Day propos'd be the 15th of May, and the Place of call'd Gunter's Quadrant. 35

of the Sun, and Ascensional Difference as before, viz. 28 gr. 50 min. this converted into Time, make 1 h. and about 55 min. and fo long did the Sun rife before six, viz. at 4 h. and 5 min. in the Morning, and consequently sets so much after fix; and in the Winter, so much as the Sun rifes after fix, so much it fets before six. Hence the Length of the Day and Night is most easily obtain'd; for having the semidiurnal Arch, or Time from Sun-Rising to Noon, 'tis but doubling those Hours and Minutes, and you have the Length of the Day; fo 7 h. 55 min. doubled, is 15 h. 50 min. for the Length of the Day, May the 15th, which substracted out of 24h. remains 8 h. 10 min. for the Length of of the Night.

36 The Use of a portable Instrument,

HARRY WERE HARRY HARRY AND HARRY HAR

SECT. VIII.

Of the Five Stars.

HE Stars that were thought convenient (by the Author) to put upon the Quadrant, are first, Ala Pegast, a Star in the Extremity of the Wing of the Pegasus, in regard it wants but min. of Time of the Beginning of Aries; and, because 'tis not always to be feen, four more were elected for each Quarter of the Ecliptick; an Oculus Tauri, or the Bull's-Eye, whose Right Ascension in time is 4 h. 20 m. then of Cor Leonis, the Lyon's Heart, whose Right Ascension is 9 h. 54 min. the next of Arcturus, whose Right Ascension is 14h. 3 min. and the last of Aquila, or the Vulture's Heart, whose Right Ascension is 19 h. 37 min. These five Stars have all of them northern Declination, and of all others some of these will be feen (if the Air be clear) every Night in the Year. Their Use follows, viz.

I. The

call'd GUNTER's Quadrant. 37

five Stars being known, to find the Hour of the Night. First, put the Bead to the Star which you intend to observe, then take his Altitude, and by that find how many Hours he is distant from Midnight, either before or after; then from the Right Ascension of the Star, substract the Right Ascension of the Star, substract the Right Ascension of the Difference; for this Difference being added to the observ'd Hour of the Star from the Meridian, or Midnight, shall shew how many Hours the Sun is gone from Noon, which is the Hour of the Night.

2. As for Example, May 15, suppose the Sun be in 4 gr. of Gemini, and I set the Bead to Arcturus, and observing his Altitude, should find him to be in the West about 52 gr. high, and the Bead to fall on the Hour-Line of 2 Asternoon, the Hour would be 11 h. 50 min. past Noon, or but 10 min. short of Midnight: For 62 gr. the Right Ascension of the Sun converted into Time, makes 4 h. 8 min. which if we take out of 14 h. 3 min. the Right Ascension of Arcturus, the Difference will be 9 h. 55 min.

38 The Use of a portable Instrument,

and this being added to 2 Hours, the obferved Distance of Arcturus from the Meridian, shews the Hour of the Night to

be 11 h. 55 min. as aforesaid.

3. Again, to make the Business more plain, suppose the 9th of July, and the Sun be then in 27 gr. of Cancer, I should set the Bead upon Oculus Tauri; and, observing his Altitude, should find him to be in the East, about 12 gr. high, and the Bead to fall on the Hour-Line of 6 before Noon, which is 18 h. past the Meridian, the Hour of the Night would be found fomewhat more than 15 min. past 2 in the Morning, which may be proved thus: The Right Afcenfion of the Sun is 119 gr. which converted into Time (by allowing 15 gr. for an Hour, and 4 min. for a gr.) it makes 7 h. 56 min. this substracted out of 4 h. 20 m. the Right Ascension of Oculus I auri (adding the Circle, or 360 gr. that Substraction may be made) the Difference will be 20 gr. 24 min. and this being added to 18 h. (which was the observ'd Distance of the Star from the Meridian) shews, that the Sun (abating 24 Hours, or the whole Circle) is 14. 11 11 11 h, 24 11 h. 24 min. past the Meridian, and therefore 24 min. past 2 in the Morning.


4. Now, if you have a Nocturnal upon the Backside of the Quadrant, you need not use this Equation of Right Ascension; for, knowing the Time of the Year, when the Star will be South at Midnight, you may bring that Time to the Hour observed; then will the Day of the Month, wherein you made your Observation, point at the Hour of the Night required.

5. As in the first Example, where, on the 15 of May, the Bead, fet to ArEturus, fell upon the Hour-Line of 2 Afternoon; because this Star will be in the South the 14th of October, compleat at Midnight, you may place the 14th of October at the Hour of 2; so the 15th of May will point to 11 h. 55 min. as before, the Hour of the Night desired.

6. And in the second Example, of the 9th of July, where the Bead, set to the Bull's Eye, fell on the Hour-Line of 6 before Noon; because the Bull's Eye will be in the South the 16th of May, compleat at Midnight, you may turn the 1.6th of May to the Hour of 6, and

D 2 thereby

40 The Use of a portable Instrument, thereby you shall find the 9th of July, to point unto 2 h. 24 min. as before.



SECT. IX.

Of the Quadrat.

divided, and the other two Sides, next unto the Center, may be supposed to be divided, each of them into 100 equal Parts; that which is next to the Horizontal Line, contains the Parts of Right Shadow; the other next to the Sights, the Parts of Contrary Shadow. Its Use follows.

2. Any Point being given, to find whether it be level with the Eye or not

If you lift up the Center of the Quadrant, so as the Thread with the Plummet may play easily by the Side of it, and look through the Sights to the Place given, and the Thread fall upon the Line AB, drawn by the Side of the Quadrant (call'd the Horizontal Line)

Line) then is the Place given level with the Eye; but if it fall within the fald Line on any of the Divisions, then it is higher; if without, it must needs be lower, than the true Level of the Eye.

3. To find any Height above the Level of the Eye, or a Distance at one Observation.

Look through the Sights to the Place, going nearer or farther from it, till the Thread fall on 100 Parts in the Quadrant; fo shall the Height of the Place, above the Level of the Eye, be equal to the Distance 'twixt the Place and the Eye. But if the Thread fall on 50 Parts of Right Shadow, the Height is but half the Distance; if it fall on 25, it is a Quarter of the Distance; if on 75, 'tis three Quarters of the Distance; for as oft as the Thread falls upon the Part of Right Shadow, it holds this Proportion, viz.

As 100 is to the Parts upon which the Thread falls: So is the Distance to the Height required.

D 3:

And

42 The Use of a portable Instrument,

And so on the contrary,

As the Parts cut by the Thread are to 100: So is the Height to the Distance.

4. But when the Thread shall fall upon the Parts of Contrary Shadow, if it fall on 50 Parts, the Height is double the Distance; if on 25, 'tis four Times as much as the Distance: For as often as the Thread falleth on the Parts of Contrary Shadow, the Proportion is,

As the Parts cut by the Thread are to 300: So is the Distance unto the Height.

And the contrary,

As 100 is unto the Parts cut by the Thread: So is the Height unto the Distance.

And what is here said of Height and Distance, the same may be understood of Height and Shadow.

5. To find a Height or Distance at two Observations.

Suppose

call'd Gunter's Quadrant. 43

Suppose the Place, which is to be measured, could not otherwise be approached, and yet it were required to find the Height and Distance; first, I make choice of a Station where the Thread may fall on 100 Parts in the Quadrat, and 45 gr. in the Quadrant; then shall the Distance from the Object be equal to its Altitude. But if I go farther, in a direct Line with the former Distance, and make choice of a second Station, where the Thread may fall on 50 Parts of Right Shadow, then shall the Distance, from the Object to my Station, be double the Height of the Object: Wherefore, if I measure the Difference between the two Stations, it will be equal both to the first Distance, and the Altitude of the Object. But if I cannot make choice of fuch Stations, then I take such as I may; as suppose one, where the Thread falls upon 50 Parts of Right Shadow, and the second upon 40, and suppose the Altitude of the Object 100, &c. the Proportion holds,

As 50 Parts are unto 100, the Side of the Quadrat: So is 100, the supposed Height, unto 200, the Distance.

And,

44 The Use of a portaInstrument,

And,

As 40 Parts, at the second Station, are unto 100: So is 100, the supposed Height, unto 250, the Distance desired.

Wherefore the Difference, between the two last Stations, should seem to be 50; and then, if, in the Measuring of it, I find it to be either more or less, the Proportion holds, as from the supposed Difference to the measured Difference; so from Height to Height, and from Distance to Distance: As if the Difference between the two last Stations, being measured, were found to be 30; then say, by the Rule of Proportion, or the Golden Rule,

As 50, the supposed Difference, is unto 30, the true: So is 100, the supposed Height, unto 60, the true Altitude.

And,

As 200, the supposed Distance, is unto 120, the true: So is 150, at the second Station, unto 250, the Distance required.

6. If

call'd Gun = R's Quadrant. 45

tude, and you would thereby know the Distance, work as you did before; only that which was then the Parts of Right Shadow, must now be the Parts of Contrary Shadow; and that which was then the Parts of Contrary Shadow, must now be the Parts of Right Shadow.

SECT. X.

The Description and Use of the Nocturnal, which is usually put upon the Backside of Gunter's Quadrant.

know the Hour of the Night by the Stars: It consists of two Parts, the Fixed and the Moveable. The fixed Part, being the outermost Circle, is divided into 24 Parts, representing the Hours of the Night; of which so many as are necessary are inserted, which are from 4 in the Asternoon till 8 in the Morning: Each Hour is divided into Halves, Quarters, and Half-Quarters, and farther.

46 The Use of a portable Instrument,

ther, according as the Instrument will bear. This Part must be fixed on the Back of some other Instrument, or it

may be used by itself.

2. The other Part is moveable on its Center; the outward Circle next the Edge is divided into 365 Parts, repre-fenting the Days of the Year; and these are divided into Months, which are distinguished by Divisions and Figures, to know every Day, as is usual, and the Name to each Month. Within this moveable Place, there is inferted five Constellations of those Stars that are next the North Pole; and in some a Star of the first Magnitude, called Hircus; all which (without incumbring the Nocturnal with more) are fufficient for this Use. There is likewise a Thread to be added, lying on the Diameter from 12 to 12, which ferves for a Meridian; to which those that please may have added a Scale of Declinations, and a Circle divided into 360 gr. for finding the Right Ascension, and Declination of these Stars; but that Work may be done with ease many other Ways, and is usually omitted in the Instrument.

2. The

call'd Gunter's Quadrant. 47

2. The Use of this Instrument is thus: First, learn to know the Stars, which are usually placed upon this Instrument, which is very easy from the bare Inspection of the Nocturnal itself, and they are most easily known in the Heavens: If you turn yourself to the North, and there observe the seven great bright Stars, in the greater Bear, called by most People Charles's Wayne, those four in a square Form are in the Body of the Bear, the three other are in the Bear's Tail: Having found thefe, take notice that the two Stars in the Square, farthermost from the Tail of the Bear, do in a streight Line point to another Star almost as big as they are (which is in the Tip of the little Bear's Tail) and called the Polar Star, or North Star, and very near, tho' not exactly, in the Pole. These being known, the rest will be easily found out, by bare Inspection of the Instrument.

3. The Stars being known, now for the more certain Finding of the Hour of the Night, it will be necessary to find a Meridian, which is a Line, or Arch, conceived to be drawn through or near the North Star, and so right over your

Head:

48 The Use of a portable Instrument,

Head; and round about by the South, just under your Feet, until it meet in the North Star again; or, to spake more properly, it is a Circle that passes through the North and South Poles, and the Zenith and Nadir of the Place, and divideth the Firmament the East and West Hemispheres: And to find this Meridian, I have already shewn in the Use of the Quadrant. One good Way may be, in some Garden, or private Place, where you may clearly fee the Stars, both above and below the Pole, to pitch exactly upright, and at some Distance, two Poles of a good length; but this must be done just at Noon, so that the Shadow, made (by the Sun at that Time) of them both, may be in a direct Line, which is a true Meridian. This being premised:

4. If at any Time of the Night you would find the Hour by the Nocturnal, stand by the Southermost of your Poles, (before set up) bring it in a Right Line with the other, so that the Pole by you may hide the other from your Eye; then see what Stars are hid from your Eye by these two Poles, either above or below the Polar or North Star, which

call'd Gunter's Quadrant. 49

are then upon the Meridian: Then coming to your Nocturnal by the Help of your Thread, bring the moveable Part thereof into the fame Polition as you find the Stars in the Firmament, fetting those upon the Meridian under the Thread, that you find to be hid from your Eye by the two Poles: Then find, in the moveable Part, the Day of the Month of your Observation, and just against it, on the fixed Part, you shall find the Hour of the Night.

Way to have a Meridian for a while to exercife, until you are well acquainted with the Operation and Performance; yet you may come very near to the Matter (and near enough for common Use) without a Meridian, thus: When you know the Polar Star, place your Body directly upon the Star, and imagining a Line drawn from the Zonith, or Point directly over your Head, through the said Star, and so directly down to the Earth, observe what Stars, of those that are on your Plate, either over or under the Pole Star, are upon that supposed Line which is your Meridian; then, coming to your Nocturnal,

50 The Use of a portable Instrument,

place the moveable Part in the same Posture, and just against the Day of the Month, on the moveable Part, is the Hour of the Night.

6. An Example or two will make

all plain.

Suppose, upon the first of August, in the Year 1731, finding, by either of the former Ways, that the two Stars of Charles's Wayne, that are farthest from the Tail of the Bear (which, as I said before, point directly to the Polar Star) are on the Meridian below the Pole; then, coming to my Nocturnal, I bring those Stars under my Thread below the Pole, and then, looking for the Day of the Month August 1st, I find that 'tis almost half an Hour past one a Clock in the Morning. And the same Night finding that Star in the Nose of the great Bear, to be on the Meridian below the Pole, I place this under my String in my Nocturnal, and feek the Day of the Month, and against it I find it to be half an Hour past 10 at Night: And so upon July the 1st, finding the Star Hircus (a Star of the first Magnitude) to be on the Meridian below the Pole, I bring my moveable Part to its proper

proper Place, as before directed) and find it to be half an Hour past nine a Clock at Night; et sic de ceteris. Usus promptus facit.

SECT. XI.

Shewing the Use of Gunter's Quadrant, in taking the Declination of a Plane, by help of a Horizonial Dial.

N the back Side of the Quadrant, you may draw a Circle, and divide it into four 90 gr. or 360 gr. Draw two Diameters, and to one of them write Axis of the Plane, and the other will cut it at the Right Angles in the Center; to the Ends of these Diameters, put sour Capital Letters, E, W, N, S, to signify East, West, North, South: On the Center of this Circle, let there be a small Pin, about two Tenths of an Inch long, on which a Horizontal Dial is to move; then, having fixt your Quadrant-Edge to the Plane.

Plane, whose Declination you are feeking, you must be very careful that it be horizontal, on which fet your Horizontal Dial, (made for the same Latitude) and on the Meridian-Line of of the Dial, let there be a small Tooth, to point at the Degree of the Circle, on the back Side of your Quadrant. These Things being thus prepared, by your Quadrant find the true Hour of the Day, to which Time fet the Horizontal Dial, that now moves upon the Center of the Circle, on the back of your Quadrant; and at the same Time will the Tooth or Index, that issueth from the Meridian Line of your Dial, cut the Circle in Degrees of the Planes Declination. This is a very plain and familier Method, of finding the Declination of a Plane, and much easier than that found by the Sun's Azimuth, till you be well acquainted with Sphericks.

Here may feem some Difficulty to the young Tyro, in finding the Hour the Dry, if he has not got an Altronomial Quadrant, and well skill'd in Sphericks: But this Obstacle is soon removed by Help of an Univerfal Ring-Dial, fuch as is described in this Ap-

pendix;

pendix; but then I advise it may be about nine Inches Diameter, for that is the best Size for this Purpose; for if it is any less, the Hours will be too small, to distinguish the Time to that Exactness, as is here required; and if the Diameter of the Dial be more, there will be too great a Penumbra attending the Ray of Light, that is cast upon the Equinoctial; therefore a Dial about nine Inches Diameter, feems to be the best for this Purpose.

ක්සියම් කම්කාලය මුතුම් කලය මුතුම් කළ කලය මුතුම් කලය මුතුම් කලය. මුතුම් මුතුම් කලය මුතුම් කලය මුතුම් මුතුම් මුත මුතුම් කලය මුතුම් ම

SECT. XII.

The Description of the Universal Ring-Dial, which sheweth the Hour of the Day in any Part of the World, and the Latitude also.

HIS Instrument is generally known, and very useful for most Persons, and a Pocket Companion. It is projected out of two great E 3

Circles of the Sphere; also an Axis, and a little Ring to hang it by; and the Making thereof is well known to the Workman. 2. The greater Circle is the grand Meridian, one Quadrant, or Quarter thereof, is divided into ninety Degrees, to set it to the Latitude of the Place wherever you are. On the other Side of the Meridian is a Quadrant of Altitude, to take the Height of the Sun, thereby to find the Latitude of the Place.

The lesser of these two Circles (though both represent great Circles of the Sphere) is the Equinoctial, divided into 24 equal Parts, or Hours, with their Halves and Quarters, which are numbred but from three in the Morning, to Nine at night; the rest of the Hours are lest out, being seldom or never used.

3. The Diameter, or broad Plate, hath a slit in the Middle, and upon one Side are the Months and Days of the Year, graduated to every fifth Day; on the other Side is the Sun's Declination, from the Equinoctial to every fifth Day, which is to be used with the

Qua-

Quadrant of Altitude, to find the Latitude of the Place. The little Ring is made to slide along the Quadrant, with a small Tooth, to set it to the Latitude: And the Use of this Portable and General Instrument is as followeth.

SECT.

SECT. XIII.

The Use of the Ring-Dial in several Examples; first to find the Latitude, and secondly the Hour of the Day in any Place throughout the habitable World; and therefore an excellent Companion for all Travellers, Gentlemen, or others, of what Degree or Quality soever: Tis fit they should accommodate themselves with both this, and the Quadrant; the one being General, the other Particular; and the one supplies the Defects of the other.

1. For the Elevation of the Pole, or the Latitude of the Place.

Piece, (or Bride, so called by some) to the Day of the Month, then turn the other Side, and right against the Hole is the Sun's Declination for that

that Day. The fame Day take the Sun's Meridian Altitude, (which is his Height at Noon every Day) and may be performed by this Instrument, thus: Put a Pin into the Hole, which you shall find in the greatest Circle; then move the Tooth to the beginning of the Degrees in the lesser Quadrant, and turn the Pin next to the Sun, and that Degree which is cut by the Shadow of the Pin, is the Sun's Altitude, (which must be taken just at Noon, as aforesaid.)

2. If the Time of your Observation be from the 10th of March, to the 13th of September, you must substract the Declination out of the Meridian Altitude of the Sun; but from the 13th of September, to the 10th of March, add, and the Sum, or Difference shall be the Complement of the Latitude, or Equinoctial Height, which being subducted out of 90 gr. leaves

the Latitude of the Place.

EXAMPLE I.

1. Suppose the Latitude was unknown, and upon the 11th Day of June; you desire to find it then, by the former Rule, find the Sun's Declination for that Day, which will be 23 gr. and 29 min. (or the Sun's greatest Northern Declination; then take the Sun's Altitude, just at 12 of the Clock, which at, and near London, will be found 61 gr. 57 m. then substract the Sun's Declination, 23 gr. 29 m. out of 61 gr. 57 m. and the Remainder will be 38 gr. 28 min. the Complement of the Latitude or Equinoctial Height, which subducted from 90 gr. leaves 51 gr. 32 m. for the Latitude of London.

EXAMPLE II.

2. As the former Directions shew how to observe the Latitude, the longest Day in the Year, when the Sun has his greatest Northern Declination; so here admit, that on the 10th of December, when the Sun's greatest Southern Declination is also 23 gr. 29 min. And the Meridian Altitude, but 14 gr. 59 m. add these two Numbers together, and they make 38 gr. 28 m. which deducted from 90 gr. leaves the Latitude as before.

2. To find the Hour of the Day. You must set the Tooth or Index, to the Height of the Pole, or Latitude of

the

the Place, and slide the Hole in the Plate to the Day of the Month; then draw out the Equinoctial, or lesser Circle, and as near as you can guess at the Hour, and turn the Hole to it; then hold the Instrument by the little Ring, and move it till the Sun shine through the Hole, upon the middle Line in the Equinoctial, and that is the true Hour of the Day desired; and the Meridian, as it hangeth, lies in the Meridian of the World, truly North and South.

AN

APPENDIX.

The Use of Gunter's Line in messuring Board and Timber.

ADmit a Board be 10 Inches broad, and 20 Foot long, I demand the Content.

Set one Foot of the Compasses in 12, and extend the other to 10; then set one Foot in 20, and extend the other back-

backwards, and it will fall in 16 Foot and a half, which is the Content of the Board required.

2. Admit a Board be 15 Inches broad, and 27 Foot long, I demand the Con-

tent.

Set one Foot in 12, and extend the other to 15; then fet one Foot in 27, and extend the other forwards, and it will fall in 34 Foot which is the Content requir'd.

3. Admit a Board to be 7 Inches and a balf broad, and 29 Foot long, I demand

the Content.

Set one Foot in 12, and extend the other to $7\frac{1}{2}$; then fet one Foot in 29, and extend the other backwards, and it will fall in 18 Foot, the Content required.

4. Admit a Board to be 27 Inches broad, and 9 Foot and a half long, I demand the

Content.

Set one Foot in 12, and extend the other to 27; then fet one Foot in 9½, and extend the other forwards, and it will fall in 21 and a Quarter, which is the Content defired.

5. Admit a Piece of Timber to be 18 Inches square, and 20 Foot long, I desire to know bow many square Feet there is

contained in this Piece of Timber.

Set one Foot in 12, and extend the other to 8, then fet one Foot in 20, and extend that Distance twice backward, and it will fall in 8 Foot 3 Quarters, which is the true Content required.

6. Admit a Piece of Timber to be 9 Inches and 2 Square, and 27 Foot long, I de-

mand the Content.

Set one Foot in 12, and extend the other to $9\frac{1}{2}$, then fet one Foot in 27, and extend that Distance twice backward, and it will be in 17 Foot, the true Content required.

7. Admit a Piece of Timber to be 18 Inches square, and 9 Foot long, I demand

the Content.

Set on Foot of the Compasses in 12, and extend the other to 18, then set one Foot in 9, and extend that Distance twice forward, and it will fall in 20 Foot and $\frac{1}{2}$, the Content desired.

8. Admit a Piece of Timber to be 22 Inches square, and 35 Footlong, I demand

the Content.

Set one Foot in 12, and extend the other to 22, then fet one Foot in 35, and extend that Distance of the Com

F paf

L ..

passes twice forward, and it will fall in 118 Foot, which is the Content required.

9. Admit I have the Breadth of a Beard in Inches, to find how much in

Length will make a Foot superficial.

Extend the Compasses from the Breadth to 12, the same will reach from 12 to the Content forward, if the Breadth be less then 12, and backward of the contrary.

复杂茶菜菜菜菜菜菜菜。菜菜、菜菜、菜菜菜菜菜菜菜菜菜菜

SECT. II.

Shewing the Use of the Carpenters Two-Foot-Rule, in the Mensuration of Board and Timber, &c.

BECAUSE cross Multiplication is better understood, and consequently imbraced by the common Artificers, than Decimal Arithmetick, I shall here subjoin a Table with Directions, for instructing those unacquainted with it, which is this:

Cross Multiplication-Table.

Factors	Feet	Inches	Parts	Seconds	
Feet	Feet	Inches	Parts	Seconds	
		by 12			
4	1	Parts			
1		by 12		1	
E .		Seconds	•	3 %	
		by 12			
Seconds		Thirds		a de la companya de l	
	by 12	by 12	by 12	by 12	

Explanation.

1. Feet multiplied by Feet produce Feet.

2. Feet by Inches produce Inches, which divided by 12 give Feet.

3. Feet multiplied by Parts of an Inch, and divided by 12, are Inches.

4. Inches multiplied by Feet, and

divided by 12, are Feet.

5. Inches multiplied by Inches, and that Product divided by 12, give Inches.

6. Inches multiplied by Parts produce Seconds, (or a Quantity less than

the Part of an Inch) which divided by 12 give Parts of an Inch.

7. Parts of an Inch multiplied by Feet, and divided by 12, give Inches.

8. Parts of an Inch, multiplied by Inches, and divided by 12, gives in the Quotient Parts of an Inch, and so on, as the Table expresseth, to Seconds, which are a Quantity 12 Times less, than the Parts of an Inch; as for Example: Let it be required to

Note. In adding the feveral Parts together, you must carry 12 in all the Denominations. ExamExample 2. Multiply me 7 Feet, 8 Inches, by 9 Inches, and 3 Parts.

Operation.

Example 3.

F 3

Example

Exam. 4. perform	4. performed			Duodecimally			
		F.	I.		Ś.		
Multiply	me	9	10	3	9		
	By	8	11	9	6		
	4	IL	I	10	6		
7	4	8	9	9			
9 0	5	5	3				
78 10	6	0					
Product. 88 6	9	I	2	7	5		

Note. You must begin with this Ex. ample at the right Hand, and carry the 12, as you do the tens in common Multiplication, and fay 6 Times 9 is 54, that is 6 and carry 4 Twelves; then 6 times 3 is 18, and 4 I carried is 22, that is ten above 12, and carry Twelve; then 8 times 10 is 80, and TI carry'd is 81, that is 1 and carry 6 Twelves; then 6 times 9 is 54, and 5 I carry'd is 59, that is 4 Feet and 11 Inches to fet down, as you fee in the Example: Then you must go on just thus with every one of the other Multipliers, which done, you must add the four several Products together, by carrying the 12, and then I find the Product to be 88 Feet, 6 Inches, 9 Parts (to 3 of an Inch) 1 Second, 2 Thirds, 7 Fourths, and

6 Fifths, which is the exact Product of F. I. P. S.

Multiply by 8 11 9 6
The like is to be observed of any other Feet and Inches whafotever.

Being thus prepared with Duodecimal Arithmetick, commonly called Cross Multiplication, you must in the next Place understand the Artificers common Two-Foot-Rule, which I thus describe. This Rule is made of dry Boxwood, two Foot long, with a strong Brass-Joint, to fold up to a Foot, for the Conveniency of the Pocket, on which there is a Line of Inches, numbred from one End to the other with Figures, 1, 2, 3, 4, &c. to 24; and each Inch is divided into 8 equal Parts, or half Quarters of an Inch; under this Line of Inches is placed the Line of Board-Measure, with the Table at the End, which is to shew how much in Length and Breadth will make a superficial Foot; and this Line is thus made, viz. As the Length or Breadth of the Board is to 1; so is 144 to the other; that is, If a Board be 8 Inches broad, how much in Length will make a Foot of that Board? Opera-

Operation.

As 8: 1:: 144:18. So 18 Inches in Length, and 8 in Breadth, is just one Foot of Board. So likewise, if the Breadth of a Board be 6 Inches, the Length to make a Foot; must be 24 Inches, or 2 Foot, and if the Breadth of the Board be 7 Inches, the Length of that, to make a Foot, must be 1 Foot, 8 Inches, and 6 Tenths of an Inch; but if the Breadth be only 5 Inches, then the Length of the Board, to make a Foot, must be 2 Feet, 4 Inches, and 8 Tenths of an Inch: As this Table, which is on one End of the Rule, more fully sheweth.

A Table of Board-Measure.

Inches	I	2	3	4	5	6	17	8	
Feet	12	6	4	3	2	2	I	I	iii na
Inches	0	0	0	0	4.8	0	8.6	6	

But when the Breadth of the Board exceeds the Table at the End of the Rule, then find it along the Side, and, right against the Inches of the Breadth, you will find, in the Line of the Board-Measure,

Measure, the Inches in Length of that Board, that will make a Foot; as, Suppose the Breadth of the Board be 10 Inches, what Length will make a Foot? Look for 10 Inch. in the Line of Inches, and right against, it in the Board Measure Line, you will find 14 Inches, and 4 Tenths of an Inch, and fo much in Length will make a Foot. Thus have I given you, the full Explanation of the Line of Board-Meafure, and now I shall shew you its Use; how expeditiously to measure any Board, which is perform'd thus: Take the Rule in your Hand, and measure the Breadth of this Board, which I will here suppose to be 11 Inches, then look on your Board Line, and fee how much in Length, and that Breadth, will require to make a Foot, which you will find there to be 13 Inches, and one Tenth Part of an Inch. Lay your Finger at 13 Inches, and begin at the End of the Board, and measure along the Side, till you come to the other End; then, as often as you find 13.1 Inches, in the Length of the Board, fo many Feet of Board is contained therein.

Example 2. Suppose you measure with the Two-Foot-Rule, and find the Breadth of a Board, &c. to be 23 Inches, and look upon the Board-Measure Line, and find the Breadth of that Board that will make a Foot, which is $6\frac{1}{4}$ Inches; here hold your Finger, or mark it with Chalk; then begin at the End, and measure along the Board, and as often as you find $6\frac{1}{4}$ in the Length, so many Feet doth that Board contain.

If your Board to be measured be more than 24 Inches broad, so that you cannot see upon the Line of Board-Measure, how many Inches in Length goes to make a Foot of that Breadth; I say, in this Case, you must have recourse to the Proportion, made use of in making the Line above given, and then proceed to measure the Board as before. Example: There is a Table, whose Breadth is 27 Inches, and Length 7 Foot, how many Feet of Board doth it contain?

First find how much of that Breadth

will make a Foot, thus,

As 27: 12:: 12: 5.33. Or, as 27: 1:: 144: 5.33. Here you fee, that 5.33

Length is a Foot. Mark your Rule with Chalk at 5.33, and measure along the Side of the Table, and you will find it to contain 15 Feet, and almost three Quarters of a Foot.

The common Artificers Rule is now much improved, by adding a Slider in one of the Legs, which, if they do but take the Pains to understand, will much facilitate the Work of Meafuring Superficies and Solids; and, for the fake of those that are unacquainted with it, I shall here give them, to understand, that the Lines, upon those Sliding Rules, were the Invention of Mr. Edmund Gunter, and so are known, by the Name of the Line of Numbers, or Gunter's Line, being at first deduced from the Logarithms, which are composed of a single and double Radius, to slide by each other. The Radius is Divided into nine unequal Parts, in a Geometrical Proportion, and figured, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, called Primes.

These Divisions are again divided into 10 like Parts, called Tenths, and those again divided (where the first

Divisions are large) into five like Parts called Centisms. And the Figure, at the beginning of the Rule, towards the left Hand, may be either .1, 1, 10, 100, or 1000, then will the same Figure, in the Middle of the Rule, (when a double Radius) be 1, 10, 100, 1000, or 10000, and at the End, to the right Hand, 10, 100, 1000, 10000, or 100000, encreasing in a tenfold Proportion; which, being well understood, will give you the full Numeration upon the Rule. The next thing to be known is, how to multiply upon the Rule, which is thus perform'd: Set I upon the Slider to the Multiplicand upon the Stock, and against the Multiplier upon the Slider, is the Product upon the Stock. Or, you may take I upon the Stock, and the Multiplicand upon the Slider, and it will produce the same Thing.

Example. Let it be required to multiply 2 Feet, and a Quarter, by

5 Feet and a Half.

Operation.

Set I upon the Slider to 5.5 upon the Stock, (or to 2.25, it matters not

not which) and against 2.25 on the Slider, is 12.375, that is, 12 Feet and a Quarter, and half a Quarter of a Foot.

Other Examples.

Products.

Multiply \(\begin{array}{c} 7.3 \\ 5.7 \\ by \\ 9.4 \end{array} \) \(\begin{array}{c} 147.5 \\ 7.6 \\ 7.6 \\ 7.6 \end{array} \) \(71.4 \\ \end{array} \)

Division on the Sliding-Rule.

The Rule.

SET 1 on the Slider, to the Divisor on the Stock-Piece, and against the Dividend on the Stock-Piece, is the Quotient on the Slider.

Example. Let it be required to di-

vide 12.375 by 2.25.

Operation.

Set 1 on the Slider, to 2.25 on the Stock-Piece, and against the Dividend 12.375 on the Stock, is 5.5 on the Slider.

Other Examples.

Divide
$$\begin{cases} 147.5 \\ 77 \\ 71.4 \end{cases}$$
 by $\begin{cases} 7.3 \\ 5.7 \\ 9.4 \end{cases}$ $\begin{cases} 20.2 \\ 13.5 \\ 7.6 \end{cases}$ Qu.

Three Numbers given, to find a fourth in a Direct Proportion.

The Rule.

As the first Term upon the Slider, is to the second on the Stock; so is the third upon the Slider, to the fourth on the Stock.

Examp. If 10 Foot of Board cost 155. what will 50 Feet cost at that Rate?

Operation.

Set 10 upon the Slider, to 15 on the Stock-Piece, and against 50 on the Slider, is 75 on the Stock, that is, 3 l. 15s. the Price of the 50 Feet of Board.

Three Numbers given, to find a fourth in a Reciprocal or Inverted Proportion.

The Rule.

As the third Term upon the Slider, is to the first upon the Stock; so is the second upon the Slider, to the fourth upon the Stock Piece.

Exam-

Example. If 6 Men can build a Wall in 9 Days, what Time will 3 Men be in doing the same.

Operation.

As 3 upon the Slider, is to 6 on the Stock; so is nine on the Slider, to 18 on the Stock; and that Time 3 Men will require to finish the same Work.

To measure Board, or Glass, &c. on the Sliding-Rule.

A S the Length or Breadth in Inches, is to 144; so is the Length or Breadth in Inches, to the Content in Feet.

Example. Suppose a Board or Pane of Glass be 16 ½ Broad, and 32 ½ long, how many superficial Feet doth it contain.

Operation.

Set 16 on the Slider, to 144 on the Stock-Piece, and against 32 on the Stock, is 3 Feet and 7 Tenths of a Foot upon the Slider, the Answer fought.

G 2

Hav-

Having the Length and Breadth of any oblong Superficies in Feet, to find the Content in Feet.

This is no more, than to multiply the Length in Feet by the Breadth in Feet, and you will have the Content in Feet.

Example. Suppose the Breadth of a Board or Glass be 2½, and the Length 17½, how many Feet doth it contain?

As 1: 2.5:: 17.75: 44.375, that is, 44 Feet, and a Quarter and Half of a Foot, the Content.

Secondly, for Timber.

There is a Line upon the Artificer's Two-Foot-Rule, called the Timber Measure; the making of this Line is thus: As the Square of the Side, of any Piece of Timber or Stone, is to i; so is the Cube of 12, viz. 1728, to the Length in Inches, that will make a Foot solid.

Example. Let the Side of a square Piece of Timber or Stone be 3! Inches, how many Inches in Length will make a solid Foot?

Operation.

First square 3-1 which is 12.25; then fay, As 12.25: 1:: 1728: 14.1, that is, 14 Feet, and one Tenth of a Foot, that Side will require to make a folid Foot. And after the same manner is the Line of Timber-Measure upon the Rule made.

The Use of the Line of Timber-Measure is this.

Suppose a Piece of Timber or Stone be 17 Inches square, at both Ends throughout, and 10 Feet long, how many folid Feet doth it contain? Look on the Line for Timber-Measure, and find how many Inches in Length will make a folid Foot, which you will find to be 5.97 Inches, which note; then begin at the End of the Timber, and measure along; for as often as you find 5.97 in the Length, so many Feet of Timber will you find contain'd therein, which in this Example is 20 Feet, and 114 Parts of a Foot. By the Sliding-Rule, two Operations at

once setting the Rule.

Say, As 12 is to the Side of the Square 17; so is 10 Feet the Length, to a fourth Num- G_3

Number 14.1. Then, without stiring the Slider, say, As 12 is to 17; so is the fourth Number 14.1, to the Content

in Feet, 20 1 24.

Example. 2. In round Timber, it is the common Way, used amongst Artificers, to girt the Tree, and to take of that Girt for the Side of the Square; which let be 7 Inches, and the Length 20 Feet, what is the Content?

Operations.

As 12:7::20:11.6.

Now fay, As 12 7:: 11.6: 6.76 Feet.

Example. 3. Let the Side-Square be $8\frac{1}{2}$, and the Length 30 Feet, how many Feet is contain'd in this?

Operation.

As 12: 8.5:: 30: 21.25

Now fay, As 12: 8.5: 21.25: 15

Also, let the Side-Square be 14, and the Length 36 Feet: Then,

 A_{\bullet}

As 12: 14: 36: 42.

Now fay, As 12: 14:: 42: 49 Feet.

Timber may be measured yet more expeditiously, than by what I have given above; for if your Rule has the Girt-Line on it, you may find the Content at one Operation. Now the Girt-Line is nothing else, but a single Line of Numbers, made to slide by the Side of a double Line of Numbers, and begins at 4, and continued to 40; this contains just one Radius of the Line, which answers to a double Line of Numbers; and the Reason of the Line being thus disposed, is because 40 Girt-Feet of Timber is a Load square. This known, girt your Timber about the Middle, a fourth Part of which is the Side of the Square; then measure the Tree in Feet, and fay, As 12 on the Girt-Line, is to the Length in Feet on the Line of Numbers; so is 4 of the Circumference on the Girt-Line, to the Content in Feet upon the Line of Numbers.

Example. Let the Circumference of a Tree be 100, one fourth of that

80

is 25 Inches, and the Length 12 Feet, how many folid Feet are there in this Timber?

Operation.

As 12: 12: 25: 52 Feet.

Another Example.

As 12: 16:: 20: 44.4 Feet.

Example 3.

As 12: 17.5: 17.5: 37.2 Feet.

N. B. But if it happen, that when 12 on the Girt Line is fet to the Length of the Tree, on the Line of Numbers, that then 4 of the Circumference fall off the Line, to remedy this, there must be a new Gauge-Point found, to supply the old Gauge-Point 12, which is thus performed, viz. Set Unity on the Line of Numbers, to 12 on the Girt-Line, and against Unity on the Line of Numbers, you'll have the new Gauge-Point 37.95 on the Girt-Line, with

with which work, as has been taught above with the Gauge-Point 12, and

you'll have the true Content.

Example. Let the fourth Part of the Girt be 17.5 (as in the last Question) and the Length 17.5 Feet, what is the Content, by the new Gauge Point on the Girt-Line?

Operation.

Ga. Point. Leng.
As 37.95: 17.5:: 17.5: 37.2 Feet.

More Examples.

G. P. Leng.
As 37.95: 10.25:: 13: 12.1 Feet.


G. P. Leng.
As 37.95: 22:: 9: 12.3 Feet.

G. P. Leng.
As 37.95: 18:: 11.5: 16.5: Feet.


G. P. Leng,
As 37.95: 48:: 14.75: 72.8 Feet.


G. P. Leng.
As 37.95: 33.5:: 20.25: 96 Feet.

THE









Art of Numbring

BY

SPEAKING-RODS,

Vulgarly termed

NEPIER's BONES,

By which

The most difficult Parts

O F

ARITHMETICK,

Multiplication, Division, and Extracting of Roots, both Square and Cube, are performed with incredible Celerity and Exactness (without any Charge to the Memory) by Addition and Substraction only.

Publish'd by W. L.

LONDON: Printed for J. WILCOX, at the Green-Dragon, in Little-Britain, MDCCXXXII.



THE

ARGUMENT

TOTHE

READER

Lord Nepeir, Baron of Merchiston in Scotland, in the Compofure of those ever to be admired Tables of his Invention, called Logarithms, finding his Calculations so laborious in long and tedious Multiplications, Divisions, and Extracting of Roots, that his Invention to him must needs render itself very unpleasant, had he not known that the Labour when finished would crown both Him and his Work. He advised with divers learned Men, studious

H

To the READER.

in the Sciences Mathematical, and to them (and amongst them) especially to Mr. Henry Briggs, who (by a learned and able Divine) was styled (and not without due Respect) our English Archimedes, to him. I say, this honourable Lord imparted his Invention, who joyning Issue with him, in this Herculean Labour, brought them to that Perfection to which they are now (to the Admiration).

tion of all Europe) arrived.

In the tedious Calculation of these Numbers, the Author sinding his Work to go on but very slowly, at length, studying out for some Help by Art, to assist him in this his noble Enterprize, thinking upon several Helps, at last (by the Blessing of God) he hapned to find out this, which I here intend to describe and shew the Use of, with some Additions and Variation form what he has himself done, in his Treatise in Latin, published and printed at Edinburgh in Scot-

The ARGUMENT

land, Anno 1617, entitled, Rab-dologiæ seu Numerationes per Virgulas. The Uses whereof I shall, in the following Tractate, endeavour to render so plain and easy, that he, that can but Add and Substract, shall be made able, in a Day's Time and less, to Multiply and Divide any great Numbers, nay, and to Extract both the Square and Cube Roots.

I have begun this Treatife with the Fabrick and Inscription of these Rods, according to the Author's Description, which being not so convenient, either for Portability or Pra-Etice, as some others which I have seen and used, I have described them (I think) in the best Manner they

possibly can be contrived.

For their Use, I am sure I have done more than hitherto I have seen done, and (if I mistake not) to as good and effectual Purpose. I do not publish it as a Novelty, neither do I attribute much in it to myself besides


H 2 the

To the READER.

the Method; for, had I not been defired, I should hardly have thought upon it: However, it being done, accept it and use it, till I direct samething else to thee, which may be more acceptable, till when, I bid thee heartily

Farewel.

CHAP.



CHAP. I.

Concerning the

Fabrick and Inscription

Of these

RODS.

N the foregoing Argument I told you, that the Author and Inventor of this kind of Instrument, of which I intend to shew the Use, called it Rabdologia, and the Word he thus defines,

Rabdologia est Ars computandi per Virgulas numeratrices: That is, Rabdologia is the Art of Counting by

numbring Rods.

 H_3

I. Of

1. Of the Fabrick of these Rods, according to the Inventor's Description of them.

These Rods may be made either of Silver, Brass, Box, Ebony, or Ivory, of which last Substance, I suppose, they were at first made, for that they are (for the most part) by all that know, or use them, called Nepeir's Bones.

But let the Matter of which they are made be what it will, their Form (according to this Description) is exactly a square Parallelepipedon, the Length being about three Inches, and the Breadth of them about one tenth Part of the Length. But the Length of these Rods are not confined to three Inches, but let the Length be what it will, the Breadth must be a tenth Part thereof; but that may be accounted a competent Breadth, that is capable of receiving two numerical Figures, for there is never upon one Rod required more to be fet on the Breadth thereof.

The Breadth of these Rods, being exactly one tenth Part of the Length thereof, when 10 of these are laid together, they do exactly make a Geo-

metrical

metrical Square, and if 20 of them be tabulated or laid together, they will make a right-angled Parallelogram, whose Length is double to its Breadth; if 30 be tabulated, the Figure will be still a Parallelogram, whose Length will be three Times the Breadth; and so if 40, four times the

Length, & sic. &c.

The Rods being thus prepared of exact Length and Breadth, let each of them be divided into 10 equal Parts, with this Proviso, that nine of the ten Parts stand in the Middle of each Rod, and the other tenth Part must be divided into two Parts; half whereof must be set at the one End, and the other Half at the other End of the same Rod. Then from Side to Side draw right Lines from Division to Division, so is your Rod divided into Squares on every Side thereof. Lastly, from Corner to Corner of every of these Squares draw a Diagonal Line, and that will divide every Square into two Triangles. The Rods being thus prepared, and lined first into Squares, and then into Triangles, they are then fit to be numbred.

4 The Art of Numbring

The Figure 1, at the beginning of the Book, shews the Form of one of these Rods lined as it ought to be.

CHAP. II.

How these Rods are to be numbred.

In the two half Squares, which are at the Ends of each Rod on every Side, there are fet one single Figure, on each Side of every Rod one, in the Division at the End thereof; so every Rod containing four Sides, ten Rods will contain 40 Sides, and so consequently will have 40 single Figures at the Ends of every of them; that is, there will be upon the ten Rods amongst them four Figures of each kind, that is, four Ones, 1111; four Two's, 2222; four Threes, 3333; four Fours, 4444; four Fives, 5555; four Sixes, 6666; four Sevens, 7777; four Eights, 8888; four Nines, 9999; four Cyphers, 0000.

And here it is to be noted, that what Figure foever it be, that standeth at the Top of the Rod alone, the Fi-

gure

gure that standeth alone on the other Side of the same Rod maketh that Figure up the number 9. As for Example: If I stands on one Side, 8 will stand on the other Side, so 2 and 7 &c. as in this Table, where

If any Side of 4 side of the 7 Rods, then 2 fame Rod.

This also is to be observed, in the Figuring of every Rod, that what Figure soever standeth alone at the Top, or superior Part of the Rod, the Figure, or Figures that stand in the two Triangles, next underneath it, is double to the Figure which standeth at the Top. And the Figures which standeth at the Top. And the Figures which stand in the next two Triangles below, that is three Times as much as the Figure above. And that in

the

Times as much as the Figure above &c. till you come to the lowest Triangles in that Rod, and then the Figure, or Figures that stand in those Triangles, are nine Times as much as the Figure which standeth at the Top of the Rod.

So if a Rod have 4 at the Top thereof, the two Triangles, which are just and next under it, have only 4 in them, which is equal to 4; in the next two Triangles below there is 8, which is double to 4; in the two Triangles below them is 1 and 2, which together make 12, which is three Times as much as the 4 at the Top; the next Triangles have in them 16, which is four Times as much; the next 20, which is five Times as much; the fixth hath 24, which is fix Timesas much; the next Triangles have in them 28, which is feven Times 4; the next hath 32, which is eight Times as much; and the last Triangles at the bottom have 36 in them, which is nine Times as much. All which is visible by Figure II, at the beginning of the Book.

And

And is evident enough by this little Tabe following, which is the Table of Multiplication, commonly called Pythagoras's Table.

	The Figures in the									1	
3	Zinch	Eighth	Seventh	Sixth	Fifth	Fourh	I hird	Second	The state of the s		
	0	0	0	0	0	0	0	0	0	0	耳
	10	000	7	0	2	4	w	N	(mm)	berd	ng
	18	16	14	12	0.1	∞	0	4	2	2	igure at
	27	24	21	81	15	12	9	6	ري 	<u>بر</u>	the
	36	22	28	12	20	16	12	×	4	4	H
	45	40	25	30	25	20	15	0	7	5	op of
- Contraction of the Contraction	54	43	42	36	30	24	200	12	0	6	
		56	49	42	35	00	21	14	7	7	ich
- orthograph at the	72	64	56	48	40	32	24	16	8	~	each Rod
berg construction outposses	00	72	63	14	\$	36	27	\propto	9	9	9
The same of	Square which are										
SATURATION OF THE PARTY OF THE	Times as much as the Figure at the Top.										
1		Commission in		-		non Lan				and a second	*

Thus have you the Fabrick, Inscription, and Numbring of these Rods, according

cording to the Inventor's Contrivance of them. He makes mention of them, and hath in his Book set the Figure of the said Ten, of one of which Ten I have given you a Scheme at the beginning of the Book, which is Figure II. I will now proceed, to give you the Description of these Rods in another more commodious Form.

CHAP. III.

A Description of these Rods, according to their best and latest Contrivance.

HE Description, which I shall here give of these Rods, varies not at all from that before delivered in the Matter of which they are made, for these may be made either in Silver, Brass, Wood, Ivory, &c. Neither do they differ in their Dividing, nor yet in their Numbring: Only, whereas my Lord Nepeir maketh them square, each Rod to contain four Sides, these are made slat, consisting

fisting each Rod but of two Sides, and contain in Length about 2 Inches $\frac{1}{10}$, and in Breadth $\frac{1}{2}$ of an Inch, and in Thickness $\frac{1}{10}$ of an Inch.

One Set of these Rods consisteth of five Pieces, and therefore hath but ten Faces or Sides; whereas those of the Lord Nepeir's consisted of 40

Plains, or Sides.

Upon one of these five Pieces (a Figure whereof is at the Beginning of the Book, noted with Figure 3) you have a Cypher at the Head of the first Piece, and 9 at the Bottom thereof. Upon the second of them, you have 1 at the Head, and 8 at the Bottom; upon the Third, you have 2 at the Head, and 7 at the Bottom; upon the Fourth, 3 at Top, and 6 at Bottom; and upon the Fifth, you have 4 at the Top, and 5 at the Bottom. Every of the two Figures, at the Top and Bottom together, make 9; as 0, and 9 is 9, 1 and 8, 2 and 7, 3 and 6, 4 and 5. And here observe, that the Figures, 98765, which stand at the Bottom of the Scheme, stand with their Heels upwards, in this manner, og 199, so do all the other Figures

under them, till you come to the double Line, which is in the Middle of the Scheme, noted with A and B, at which Line, if the Scheme were cut into two Pieces, and folded or pasted on the Backside of the other Half, so that the 9 at the Bottom were placed upon the Cypher at the Top, and so 8 upon 1, 7 upon 2, 6 upon 3, and 5 upon 4; and then the Scheme cut again into five little Slippets, by the down-right Lines, these five Slippets would exactly represent one set of these Rods; for upon one of these Pieces, you should have a Cypher upon one Side, and 9 on the other; upon the next I and 8, upon another 2 and 7, on another 3 and 6, and on the other 5 and 4; both the Figures on either Side making 9, as before was defcribed.

These five Slippets do now contain the whole Table of Pythagoras before mentioned, but so few are not of sufficient Use, neither are the Ten before mentioned of the Lord Nepier's Order; for their can be but four Figures of one kind, which in all Cases is not sufficient.

There-

Therefore as these Rods are made now-a-days, they do commonly make six Sets of them, that is, 30 Pieces, which contain 60 Faces, and these will be of good Use, and there will seldom be found a Want, which, in those of the Inventor's, there will often be; except you have a great Quantity, which will be far more cumbersome, than these here described; for there is required as much Metal, or Wood, in one of his, as in four of these; and then, for his four Sides, we have here eight.

Concerning a Case for these Rods.

For the orderly keeping, and ready finding of these Rods, I have often (for myself and others) had a Box made of Walnut-Tree, or Pear-Tree, with five Partitions in it, each Partition to hold five or six Sets of these Rods, or more, if more Rods were required. Every of these Partitions being sigured on the Side thereof next the Eye, with such Figures as the Rods in such a Partition had Figures at the Top, so that the Party, that was to use them, could take them as readily out of his Partition,

tion, as a Printer can take his Letter's out of his respective Boxes to make any Word,

In this Box there is also convenient Room made for one other Rod, double in Breadth to these here described; but of the same Length and Thickness; upon the one Side whereof there is a Table or Plate useful in the Extracting of the Square-Root, and on the other Side, another for the Extracting of the Cube-Root, the Figure whereof is at the beginning of the Book, noted with Square, Cube.

But I shall forbear to say any thing of them, till I come to shew you how to extract the Square and Cube Roots, by the Help of them and the Rods.

Of a Board with a Frame, upon which to lay your Rods, when any Operation is to be wrought by them, known by the Name of a TABULAT.

In the using of these Rods care is to be had, first, of the orderly laying of them; and then, fecondly, for the keeping of them in that Polition till your Work be ended. For the effecting whereof, both neatly and certainly, there there is a little Table or Frame contrived, containing in Breadth $\frac{1}{2}$, of an Inch more than the Length of the Rods, and in Length at Pleasure, but it may well be about once and a half the Length of the Breadth.

It ought to be made of a thin Piece of Pear, or Walnut-Tree, or of such Matter as your Box or Case is made of; and it may very commodiously be contrived to be put into the Box, as I ever had them made to do, for that I found it inconvenient to carry loose.

Upon the Superficies of this Board, close to one of the Edges thereof, must be glewed, or otherwise fastned with Pins, a small Piece of the same Matter, and also of the same Length, Breadth, and Thickness of one of your Rods, which must be divided into 9 equal Parts, and Lines drawn cross the Piece; so will there be 9 Squares, in which you mult grave or stamp the nine Digits, beginning with 1 at the Top, and so descending by 2 3 4 to 9 at the Bottom thereof: And it were necessary, that these Figures (as also those which are at the Head of every of your Rods) were graven or stamped of I 3 fomefomething a bigger Figure, than the

other Figures of your Rods are.

Under the End of this Ledge, beginning at the Figures, and so continuing the whole Length of the Board, must another Ledge, of the same Matter and Thickness as the other, be glewed or pined, and then is your Tabulat sinished. A Figure whereof you have at the beginning of the Book, noted with Figure 4. It is called a Tabulat, for that, when the Rods are laid thereon, for any Operation to be wrought by them, we usually say, the Rods are tabulated.

Being thus prepared with Rods and Tabulat, you are ready for the Work intended by them, and for which chief-

ly they were invented.

Thus much for the Fabrick, Inscription, and Numbring of these Rods; let us now come to shew the

Uses of them.

CHAP. IV.

To what Use these Rods generally serve.

HE chief Uses, to which these small Rods serve, I in part intimated at the Beginning, to which Effect I shall repeat it again-For by them all Manner of Multiplications and Divisions, as also of the Extraction of both the Roots, either Square or Cube, are fo facilly and expeditiously per-formed, and that by the Help of Addition and Substraction only, that it is (as I may well fay) inconceiveable; for here is no Charge at all required of the Memory, and you shall affuredly take your Quotient-Figure in Division always certain, neither too great, nor too little, an Inconvenience so prejudicial, that I leave it to the Censure of such as have found it, to their great Loss of Time, and other Vexation which it hath put them to. But, ceasing to fay more of their Properties, I will now come CHAP. to shew their Use.

. સ્ટુલ્સ્ટ્રિક્સ્ટ્રિક સ્ટુલ્સ્ટ્રિક સેંગ્રેનિક સેંગ્રેનિક સેંગ્રેનિક સેંગ્રેનિક સેંગ્રેનિક સંગ્રેનિક સેંગ્રેનિક

CHAP. V.

How to apply, or lay down, any Numbers by the Rods.

PROP. I.

Any Number being given, how to tabulate, or lay down the same by the Rods.

ET it be required to tabulate, or lay down this Number, 3496.

First, From among your Sets of Rods (or out of your Case) take four of them, of which let one of them have the Figure 3 at the Top thereof, and lay it upon your Tabulat, close to the Edge thereof. Then,
Secondly, Take another Rod from

Secondly, Take another Rod from your Case, which hath the Figure 4 at the Top of it, and lay that also upon your Tabulat, close by the Side of the

other.

Thirdly, Take another Rod, which hath the Figure 9 at the Top of it, and lay

lay that upon your Tabulat, close by the other two.

And, lastly, take a fourth Rod, having the Figure 6 at the Head thereof, and lay that also upon your Tabulat,

close by the rest.

These four Rods thus taken, and laid upon the Tabulat, you shall see in the uppermost Row (which standeth against the Figure 1, on the Side of your Tabulat) these four Figures, 3496, that is 3496, equal to your given Number. In the second Row (against the Figure 2 of your Tabulat) you shall find the Double thereof. In the Third (against the Figure 3) you shall find the Triple thereof. In the Fourth, the Quadruple thereof. In the Fifth, the Quintuple; and so on to the Ninth and Last, in which you shall find the Noncuple of the Number given

PROP: II:

How these Rods will appear when tabulated, and, being tabulated, how to read the Multiplication; of that Number so tabulated, by any of the nine Digits.

HE four Rods being tabulated, according to the Precepts delivered in the preceding Proposition, they will appear exactly as they are reprefented in Figure 4 at the Beginning of the Book, which Figure lively represents the four Rods lying upon the Tabulat, which mind well; for upon the true Tabulating, and right Reading of the Rods so tabulated, depend the whole Work.

The Rods thus tabulated, and as you fee them in Figure 4, do to the Eye appear in the Form of a Glass-Window, every Pane thereof representing a Rhomboiades, or Diamond-Form: In the Reading of the Figures which are in these several Rhomboides's, or Diamond-Forms, observe these sew Directory.

tions

tions following, which will fully illufirate the whole Business intended, and therefore especially to be minded.

Note I. That the Figures upon the Rods are to be read beginning at the right Hand, and reading towards the left; which is contrary to our common Courseof Reading and Writing, which is from the left Hand towards the right.

II. That in every Rhomboides or Diamond, there are either one Figure, or two Figures, but never more than two.

III. If there be but one Figure in a Rhombus, then that Figure is the Figure to be set down alone, (be it either a Figure, or a Cypher) but if there be two Figures in a Rhomboides (as for the most part there is) then add them two Figures together, and set down their Sum in one Figure.

IV. But if the Sum of the two Figures, in one Rhomboides or Diamond, do exceed ten, then you must set down the overplus above ten, and keep one in Mind.

Mind, which one you must carry to the next Rhomboides.

V. That the first towards your right Hand, and the last towards your left Hand, are but half Rhomboides's or Diamonds, and never have in them more than one Figure; but all between them are whole ones, and for the most part have two Figures in them.

VI. If in either Rhomboides, or half Rhomboides, you find no Figures, but Cyphers, you must not neglect, but set them down as if they were Figures.

These Rules being rightly understood, all that follows will be familiar and easy, and these I shall explain by Example following.

Example.

For the Illustration of the preceding Rules, we will make use of those Rods which were before tabulated, therefore have recourse to Figure 4 at the beginning of the Book, where this Number 3496 is tabulated.

The Figures at the Top of the four Rods are these, 3 4 9 6, which signify

the

the former given Number 3496, and this Number stands against the Figure 1 on the Side of the Tabulat. Then, I say, that the Figures in the next Row, standing against the Figure 2 of the Tabulat, are double thereunto, which I prove thus:

Repair to the Rods as they lie upon the Tabulat, and in that Row which lieth against the Figure 2, you shall find in the first half Rhomboides, towards your right Hand (where by Rule 1 you must begin) the Figure 2; wherefore fet down with your Pen upon Paper the Figure 2; in the next Rhomboi-des, in the same Row, you will find 8 and 1, which added make 9, set down 9 on the left Hand of 2; in the next Rhombus you shall find 8 and 1 again, which is 9 also; set down 9 on the left Hand of the other; and in the last Rhomboides you shall find only 6, wherefore set down 6 on the left Hand of 9: So have you in all 6992, which is double to 3496.

Again, the Figures in the Row, which stand against the Figure 3 in the Tabulat, are triple to 3496; for in the first half Rhomboides, towards your

K right

right Hand, you have 8, set down 8; in the next Rhombus you have 7 and 1, which is 8, fet down 8 again; in the next you have 2 and 2, which is 4, set down 4; in the next Rhombus you have 9 and 1, which makes 10, let down o and carry 1; but it is the last Rhombus, and because there is never another to carry the 1 unto, you must therefore set it down: So have you this Number 10488, which is triple to 3496.

Again, the Figures standing against 4 in the Tabulat are Quadruple to 3496; for in the half Rhombus you have 4, fet it down; in the next 6 and 2, which is 8, fet that down; in the next 6 and 3, which is 9, set that down; in the next 2 and 1, which is three, set that down; and in the last half Rhomb. you have 1, which also set down: So have you 13984, which is quadruple to 3496.

Also, of the Figures against 5 in the Tabulat, the first is a Cypher, therefore put down o; the next is 5 and 3, which is 8, fet down 8; the next is o and 4, set down 4; the next is 5 and 2, that is 7, set down 7; and the last is 1,

there-

therefore set it down: So have you in all 17480, which is Quintuple to 3496.

Against 6 in the Tabulat, you have in the first place 6, set it down; then in the next 4 and 3, that is 7, fet that down; in the next 4 and 5, that is 9, fet 9 down; in the next you have 8 and 2, that is 10, set down o and carry I to the next Rhomb. where you find only 1, to which add the 1, which you carried from the Rhomb. before, and it makes 2, set down 2: So have you

20976, which is fix times 3496.

Against 7 in the Tabulat you have first 2; set it down; then 3 and 4, which is 7, fet 7 down; in the next, 8 and 6, which is 14, which being above 10, fet down 4, and carry 1 to the next Rhomb. where you have 2 and 1, which is 3, and I which you carried makes 4, fet down 4; then in the last Place you have only 2, which fet down: So have you in all 24472, which is septuple to 3496, or feven times as much.

Against 8 in the Tabulat you have first 8, which set down; then 2 and 4, which is 6, set 6 down; then 2 and 7, which is 9, fet 9 down; then 4 and 3, which is 7, fet 7 down; and lastly 2,

K 2

fet that down: So have you 27968, which is octuple to 3496, or eight times as much.

Lastly, against 9 in the Tabulat, you have in the first Place 4, set that down; in the next you have 1 and 5, which is 6, set 6 down; in the next place you have 6 and 8, which is 14, set down 4, and carry 1 to the next Rhomb. where you find 7 and 3, that is 10, which with 1 that you carried makes 11, set down 1, and carry 1 to the next Rhomb. where you find only 2, and the 1 carried makes 3, therefore set down 3: And so you have 31464, which is noncuple to 3496, or nine times as much as the tabulated Number.

Thus have I given you Examples, in shewing you how the Numbers upon the Rods are to be read and written down; and in the Delivery of this Example, I have made the whole Work, which is to follow, so plain and easy, that the meanest Capacity (I think) if he can but tell his Figures, and add any two Figures together, he may, by this here delivered, read or write down any Number that can be tabulated: And,

that you may thoroughly understand this Chapter before you proceed further, I will give you the Products of 7009078, multiplied by all the nine Digits, which I would have yourself to tabulate, and see if you find your Working by your Rods to agree with those which are here written, which Numbers if they do, you need not scruple at the most difficult that can be proposed to you; therefore study it, and try it.

7009078

Thus have I sufficiently described these Rods, and the Manner of Numbring upon them; and now I think it Time to apply them to that Use for which they were intended; namely, the more difficult Parts of Arithmetick, as

K 3

Mui-

Multiplication, Division, and Extraction of Roots; but first let me give you

An Admonition concerning Addition and Substraction.

Whereas it was the difficult Operations of Arithmetick, which, by the Benesit of these Rods, the Inventor chiefly aimed at (of which Kind he esteemed Multiplication, Division, and Extraction of the Square and Cube Roots) he omitted to fay any Thing concerning Addition and Substraction, as Things obvious to every Tyro: He therefore omitting them, begins to shew the Use of his Rods in Multiplication, whose Method I shall here follow.

遊遊旅遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊遊

CHAP. VI.

Multiplication by the Rods.

I N multiplying by the Rods, you are to consider (as in Vulgar Arithmetick) three Terms, Things, or Numbers, viz.

1. The Multiplicand, which is the

Number to be multiplied.

2. The

2. The Multiplier, which is the Number by which the Multiplicand is multiplied.

3. The Product, which is the Sum produced by the Multiplying of the

two former together.

And here note, That the Product doth contain the Multiplicand, so many Times as there be Unites in the Multiplier.

Thus for the Definition of Multiplication: Now for the Working thereof

by the Rods; for which this is

THE RULE.

First, Set down upon your Paper the Multiplicand, and orderly under it the Multiplier. [It matters not greatly which of the two given Numbers be made Multiplicand or Multiplier; but it is usual and best to make the greatest Number Multiplicand, and the leffer Multiplier. Then draw a Line with your Pen under them, and having tabulated your Multiplicand (or greater Number) look what Numbers in your Rods stand against the first Figure towards your right Hand, and that Number which you iball find upon your Rods standing against ibat first Figure found in your Tabulat, 106

set down under your Line, which you formerly drew under your Multiplicand and Multiplier; and having so done with the first Figure of your Multiplier, do so with the rest, setting them down one under another, removing every Figure one Place more towards the left Hand, than that which went before it, as is done in common Multiplication, and as you see in the following Example.

Example 1. Let it be required to multiply 3496, by 489. As if it were required to know how much 489 Times

3496 would amount unto.

First, Set down your given Numbers 3496, and 489, one under another, and draw your Line under them, as here you fee done.

Secondly, 3496, your Multiplicand, being tabulated, and 9 being the first

3496 Multiplicand | Figure to 389 Multiplier

31464 27968

13984

1709544 Product. Sec.

the right Hand in your Multiplier, look upon your Rods, what Sum

there

there stands against 9 in the Side of your Tabulat, and you shall find (as by the Rules in the second Prop. of the sifth Chap. you was directed) 31464, which is the Product of 3496, multiplied by 9; wherefore set down this Number 31464 under your Line, as you see in the Example.

Thirdly, Look what Sum upon the Rods stands against 8, which is the second Figure of your Multiplier, and you shall find 27968; set this Number under the former, moving it one Place

forward towards the left Hand.

Fourthly, Look what Sum upon the Rods stands against 4, which is the third Figure in your Multiplier, and you shall find 13984, which set down under the other, one Place more to the left Hand.

Lastly, Under these three Sums draw a Line, and add the three Sums together, and they make 1709544, which is the Product of 3496, multiplied by 489; and this 1709544, the Product, contains 3496, the Multiplicand, 489 Times.

Practife well this first Example, and compare it with the Rods as they are tabulated in Figue 4, at the beginning

of the Book, as also with the Rules in the fifth Chapter, and you may perform any Multiplication. However, I will give you one or two more Examples, and some other Ways of Multiplication.

Example 2. Let it be required to mul-

tiply the same Sum 3496 by 261.

Set the Numbers down as 3496 here is done, then look upon 3496 the Rods for the Product of 20976 3496 by 1, and you shall find it to be the same, where-6992 fore set down 3496 under the Line. Then look upon the Rods for the Product of 3496 by 6, and you shall find it to be 20976, which fet down under the other Number, one Place more towards the left Hand. Again, look in the Rods for the Product of 3496 multiplied by 2, and you shall find it to be 6992, which set down under the other two.

Lastly, Draw a Line under them, and add the three Numbers together in order as they stand, and the Sum of them will be 912456, which is the Product of 3496, multiplied by 261.

Fixann-

under

Example 3. Let it be required to multiply the same Number 3496 by 520.

Set down your Numbers as here you

fee done: — Then, because the first Fi-3496 Igure of your Multiplier, to-520 wards your right Hand, is a Cypher, wholly omit it, and multiply 3496 by 52 only, for shall you find the Product of 3496 by 2, to be 6992, which fet down: Also the Product by 5 will be 17480, which fet down under the other one Place further; then draw a Line—, and add thefe two Sums together, and they make 181792, to the which, if you add a Cypher for the Cypher which you omitted in your Multiplier, the Sum will be 1817920, which is the Product of 3496 by 520.

Example 4. Let it be required to mul-

tiply the same 3496 by 7003.

Set down your Numbers as before, and as you see here done; then, having 3496 tabulated 3496, see what the 7003 Product thereof is upon the Rods, being multiplied by 3, the first Figure in your Multiplier, and you shall find it to be 10488, which set down

under the Line: - Then the two next Places of your Multiplier being Cyphers, make two Pricks under the former Number, one under 8, the other under 4, as you see in the Example, or instead of 2 Pricks, you may make two Cyphers: - Then look on the Rods for the Product of 3496 by 7, and you shall find it to be 24472, which set down under the other Sum, beginning your Number at the fourth Place, or beyond the two Pricks or Cyphers. Lastly, Draw a Line, and add these two Sums together, and their Sum is 24482488, which is the Product of 3496 multiplied by 7003.

Thus have you four Examples in Multiplication, in which are included all the Varieties that may at any Time happen in that Rule, viz. Two, where the Multiplier confisted all of Figures, as in the first and second Example they did. — Another, where the latter Place of a Multiplier confisted of a Cypher.

—— And this last Example, where Cyphers were intermixed among the

Figures.

And thus much for this Kind of Multiplication; but, before I leave, I will shew you Ang-

Another Form of Multiplication.

Whereas, in the foregoing Form of Multiplication, which is the best and most usual, (only I insert this following for Variety) you began (your Rods being tabulated) with that Figure of your Multiplier, which stands next your right Hand; but there is no necessity for that, for you may begin with that Figure which standeth next to your lest Hand, and by so doing, and placing your several Products one Place more to the right Hand, as you did before place them to the lest Hand, those Products added together in the Form they then stand, shall produce a Product equal to the former.

Example. For our Example we will take the first Example foregoing, at the beginning of this Chapter, where it was required to multiply 3496 by 489. Set the Numbers down as before in that first Example, and as you see here done

3496 |Then 3496 being tabulated, look upon your Rods for the Product thereof, multiplied by 4, (which is the first Figure of your Multiplier towards your left Hand) and you shall find the Product thereof to be 13984, which set down. - Secondly, look the Product of 3496 by 8 (your fecond Figure) and you shall find it to be 27968, which must not be set down as in the other first Example, but as you fee it in this; 8, the first Figure thereof, must be set one Place forward towards the right Hand, as in the other, it was fet a Place backward towards the left. - Laftly, feek in your Rods for the Product of 3496 by 9, your last Figure, and you shall find it to be 31464, which set under the other two Numbers, yet one Place more to the right Hand. - So a Line being drawn under, and these three Numbers added together, produce 1709544, equal to that in the first Example: And that you may the better see the Difference of the Work, I have fet them one by the other. As

As in this
Example,
-
3496
. 489
13984
27968
31464
1709544

One Example more in Multiplication (which shall be for Advertisement and Direction) I will give, and so con-

clude Multiplication.

I said, in the general Rule for working of Multiplication (at the beginning of this Chapter) that it mattered not which of your Numbers were made the Multiplicand, or which the Multiplier, of which I will here give you a President, where the lesser Number shall be tabulated, and the greater Number only set down; and I will work it here according to this last Way of Multiplication, and the Example shall be as followeth.

Example. Let it be required to multiply 868437 by 3496, and let 3496 (the lesser Number) be tabulated.

 L_2

Let

Let the Numbers be fet as you here see, then 3496 being tabulated, begin with the first Figure towards the left 3496 Hand of your Multiplier, which here is 8, and 868437 upon your Rods find the 27968 Product of 3496 multi-20976 plied by 8, which is 27968 27968, fet that down under the Line.—Then 13984

10488

find the Product of 3496 by 6, the second Figure of your Multiplier, and

you shall find that to be 20976, this Number under the former, one Place more towards the right Hand. - Again, the third Figure of your Multiplier is 8, whose Product is 27968, as before, fet that under the other, still one Place more to the right Hand. ---In this Manner do with the other Figures of the Multiplier; as 4, the next Figure, whose Product is 13984, which also set down a Place forward. - So also the Product of 3, which is 10488, which fet down. - And lastly, of 7, which is 24472. — All these Products being fet down in the Order as you see them in the Margent, if you add

add them together, the Sum of them will be 3036055752, which is the Product of 3496, multiplied by 868437, the lesser Number being tabulated.

Other Ways of Multiplication I could have added, but these I esteem

sufficient.

CHAP. VII.

Division by the Rods.

A s in Multiplication, so in Divifion, there are three Numbers, Terms, or Things required, viz.

1. The Dividend, or Number to

be divided.

2. The Divisor, or Number by which

the Dividend is divided; and,

3. The Quotient, which is the Number issuing from the Dividend's being divided by the Divisor; and this Quotient doth always consist of so many Unites, as the Divisor is Times contained in the Dividend.

Thus much for the Definition of Division, now let us come to the Practice

L 3

38 The Art of Numbring

Of it by the Rods, to perform which, his is

THE RULE.

Tabulate the Divisor, (which is always the leffer Number of the two given) and set down the Dividend, and set the Divisor on the left Hand, and draw a crooked Line on the right Hand for your Quotient, as in common Arithmetick; then look upon your tabulated Rods (always) for the Number, less than the Number in the first Figures of your Dividend, and what Figure stands against that Number, on the Edge of your Tabulat, must be the Figure you must put in your Quotient; and that Number you must always substract from the Figures of your Dividend, and to the Remainder add another Figure, so proceeding from Figure to Figure, till your Division be wholly ended.

Example. Let it be required to divide 709544, by 3496. Having tabulated 3496, fet down your Dividend, your Dividend thereof, and a crooked Line for the Quotient on the right Hand thereof, as by the Rule preceeding you were directed, and as

you fee done in the Example adjoin-

ing.

And because, at your first setting down of your Divisor 3496, it would reach (if it were set under your Dividend 1709544) as far as the Figure 5, therefore under the Figure 5 make a Prick, to intimate how far you are gone on in your Work, and under this Prick, draw a Line quite under your Dividend, then is your Sum set down ready for Work, and will appear as you see here,

3496) 1709544 (

Your Sum thus prepared, ask how often can have you 3496 in 17095? look in your tabulated Rods for 17095, which you cannot there find, but the nearest Number thereunto among the Rods, which is less than 17095 (for you must always take a less Number) is 13984, which Number stands against the Figure 4 in the Tabulat, wherefore set 4 in your Quotient, and 13984 under the Line, and substract 13984 from 17095, and there will remain 3111;

40 The Art of Numbring

so is the first Part of your Division ended, and your Work will stand thus,

Then make another Prick under 4, the next Figure of your Dividend, so will the remaining Number be 31114:

Then look among the Rods for the Number 31114 (or the nearest less than it) and the nearest less you shall find to be 27968, which stands against 8 in your Tabulat, put 8 in your Quotient, and set 27968 under 31114, and substract 27968 from 31114, so will there remain 3146, which set over Head; so is she second Part of your Division ended, and your Work will appear thus,

3146 3111 3496) 1709544 (48

> 13984 27968

Lastly, Make another Prick under the next Figure of your Dividend, which is 4 also, making the remaining Number to be 31464, seek among your tabulated Rods for this Number (or the nearest less) but looking, you shall find the very Number, against which stands on your Tabulat the Figure 9, set 9 in the Quotient, and the Number 31464 under the Line, and substract it from 31464 the Remainder, which stands above the Line, and nothing remains; and being there is never another Figure in your Dividend, your Division is ended, and your Work will stand thus; and 3496 is contained in 1709544 489 times.

00000 3146 Divisor 3111 Quotiene 3496) 1709544 (489

> 13984 27968 31464.

2. The Art of Numbring

Another Example, and by another Way

Let it be required to divide 912456 by 3496, set down your Dividend and Divisor, draw a crooked Line for your Quotient, and also make a Prick under the fourth Figure of your Dividend, and draw a Line under your Dividend; so is your Sum prepared to be divided, and will stand thus,

3496) 912456 (

Then your Divisor 3496 being tabulated, look amongst your Rods for the nearest Number to 9124, which is less, and you shall find it to be 6992, against which stands on your Tabulat the Figure 2; set 2 in the Quotient, and this Number under the Line, and substract it from 9124, and there will remain 2132, to which Number add the next Figure of your Dividend, namely 5, and it makes 21325; under which Number draw a Line, then will your Sum stand thus,

3496) 912456 (2

6992 21325

Then among your Rods seek the nearest Number to 21325, and you shall find 20976 to be the nearest Number less, against which in your Tabulat stands 6; set 6 in the Quotient, and 20976 under the Line, substracting it from 21325, which, when you have done, there will remain 349; to 349 add the next Figure in your Dividend, which is 6, your last Figure, and it makes 3496, under which draw a Line, and your Work will stand as you see here,

3496) 912456 (26

6992 21325

20976 3496.

44 The Art of Numbring

This done, look amongst your Rods for the nearest Number to 3496, and you shall find the exact Number at the Top of the Rods, against which stand, the Figure 1 on the Tabulat; set 1 in the Quotient, and substract 3496 from 3496, the Remainder is nothing; and so is your Division ended, the Work standing thus, and 3496 the Divisor, is contained in 912456 the Dividend, 161 times.

3496) 912456 (361

A third Example ready wrought by the last and best Way of Division.

I will only fet it down ready wrought, leaving the Practice of it to yourfelf.

Let

Let it be required to divide 73020506 by 3496.

3496) 73020506 (20886 3050

6992

31005 27968

30370

-/300

24026 20976

Rem. 3050

This Sum thus divided produceth in the Quotient 20886, and 3050 remaining; so that the Quotient with Fraction and all is

20886 2010, which shews

that 3496, the Divisor, is contained in 73020506, the Dividend, 20886 times, and 3050 remaining.

This

This Example well practised, together with them foregoing, are sufficient Instructions for any Studient whatever; and be that can perform these, need not despair the most difficult that can be proposed. And fo I conclude with Division.

CHAP. VIII.

Goncerning the Rule of Three, or Golden Rule, both Direct and Reverse, or Reciprocal.

O discourse of this Rule at large, were to run into a Labyrinth; for it was the Performance of Working Multiplication and Division by the Rods that was here aimed at, and he that: can multiply and divide, may command this Golden Rule: Wherefore I will! shew you the Nature or Order of Placing the Numbers, and also the Manner of Working an Example in either of them.

The Rule of Three is that Rule which teacheth, by having three Numbers, im Proportion one to another given, too

find a fourth, which shall be in Pro-

portion to them also.

In this Rule direct, the fourth Number which is fought, is to have the same Proportion to the third, as the fecond Number hath to the first: As if the three Numbers given were 2-4 —and 8; fay, as 2 is to 4, fo is 8 to what? Multiply 4 by 8 (that is, the fecond Number by the third) and the Product will be 32, which divide by 2 (the first Number) the Quotient will be 16, which is the fourth Number in Proportion to the third, as the fecond is to the first; for as 4 the second Number, contains 2 the first Number twice; so 16 the fourth Number, contains 8 the third Number twice also.

But in the Reciprocal Rule of Three, there the Proportion is not as the first to the second, so the third to the sourth:
But as the first is to the third, so is the second to the fourth. As if the Numbers were 3, 4, and 6, say, as 3 the first Number, is to 6 the third Number, so is 4 the second Number, to what? Multiply 4 the second Number, by 3 the first Number, the Product is 12, which divide by 6 the third Number.

48 The Art of Numbring

ber, and the Quotient will be 2; for as 6 the third Number, contains 3 the first Number twice, so 4 the second Number, contains 2 the sourth Number twice also: And in this consists the Difference between the Direct and Reciprocal Rule of Three.

Questions in each Rule: And 1. In the Direct Rule.

If four Men eat two Pecks of Corn in one Week, how many Pecks will serve an hundred Men the same Time?

Men Pecks Men

Multiply 2 the second Number by 100 the third Number, the Product will be 200, which divide by 4 the first Number, and the Quotient will be 50, and so many Pecks will suffice 100 Men the same Time.

2. In the Reciprocal.

If twelve Men do any Piece of Work in 8 Days, how many Men must be imployed to do the same Piece of Work in 2 Days.

Days

Days Men Days

Multiply 8 the first Number, by 12 the second, their Product is 96, which divide by 2 the third Number, the Quotient will be 48, and so many Men will do the same Work in 2 Days; for as 8 Days are to 2 Days, fo are 12 Men to 48 Men, &c.

激素素素素等等,等等,等等等等等等等等等等等等等。

CHAP. IX.

Of the Extraction of Roots.

HE Extraction of Roots, which is the difficultest Part of Multiplication and Division, is expeditiously and certainly performed by the Rods; for the easy and expedite Performance of which, there are two Rods on purpose, one for the Square, the other for the Cube-Root, of which I will speak; first, of their Fabrick; secondly, of their Use.

Of the Fabrick of the Rods for Extract-

ing of Roots.

Of the same Matter, and of the same Length and Thickness of your other Rods, let there be made another Rod, but three Times the Breadth of the former; the Inscription on one Side ferving to extract the Square, and that on the other Side for the Cube-Root, each of which are divided into three Rows or Columns.

That which serveth for the Square-Root hath in the Top, or uppermost Square between the Diagonal thereof, these Figures 0-1, in the second 0-4, in the third 0-9, in the fourth 1-6, in the fifth 2-5, in the fixth 3-6, in the seventh 4-9, in the eighth 6-4, and in the ninth or lowermost 8-1, which are the Square Numbers belonging to the nine Digits.

In the fecond Column of the fame Rod, in the first Square is inscribed 2, in the fecond 4, in the third 6, in the fourth 8, in the fifth 10, in the fixth 12, in the feventh 14, in the eighth 16,

and in the ninth 18.

In the last or third Column, there are the nine Digits orderly descended; namely, Rod thus made is fitted for the Square-Root.

That which serveth for the Cube-Root hath in the Top, or uppermost Square of the first Column towards the left Hand, between the Diagonal thereof, these Figures, 0-01, in the second 0-08, in the third 0-27, in the fourth 0-64, in the fifth 1-25, in the sixth 2-16, in the seventh 3-43, in the eighth 5-12, and in the ninth 7-29, which are Cube Numbers orderly descending. — The second Column of this Rod contains these Square Numbers, 1, 4, 9, 16, 25, 36, 49, 64, 81, orderly descending. — The third and last Column of this Rod hath in it the nine Digits, 1, 2, 3, 4, 5, 6, 7, 8, 9, orderly descending also.

This Rod thus prepared and inscribed is fit for Extracting of the Square and Cube-Roots, a Figure of either Side whereof you have at the Beginning of the Book: That which serveth for the Square-Root, having the Word Square written over Head; that for the Cube-Root, hath Cube written over Head.

Thus having given you the Fabrick and Inscription of the Rods, I will now shew you their Use; and first,

Con-

Concerning the Extraction of the Square-Root.

In Extracting of the Square-Root, you must, as in common Arithmetick, when you have set down your Number, make a Prick under the first Figure towards your right Hand, and so successively under every second Figure; then under those Pricks draw two Lines parallel, whereinto set the Figures of your Root as you find them: Your Number being thus placed and pricked as before is directed, and as in the sollowing Example you see done, you may proceed to extract the Root thereof as followeth.

Example 1. Let it be required to find the Square-Root of this Number 12418576; first, make a Prick under 6, another under 5, another under 1, and another under 2, under which Points draw two Lines, in which you must place your Root, and then will your Number stand thus,

12418576

Take the Rod for Extracting of the Square Root, and look in the first Row

or Column thereof, for the nearest Number you can find there less than 12, (which is as far as the first Prick in your Number reaches) and you shall find 9, against which in the third Column you shall find three, set 3 under the first Point between the Lines, and 9 under the Line; and, substracting 9 from 12, there will remain 3, which fet over 12, so will your Number stand thus:

Then in the middle Co-12418576 lumn of your Rod, between 9 and 3, there stands 6; take therefore one of your-Rods which hath 6 at the Top there-

of, and lay it upon your Tabulat by. the left Side of your Square Rod, then being there is 341 to the next Prick, feek the nearest Number 1 s upon your two Rods, and you shall find the next less to be 325, against which, in the last Column of your Square-Rod, stands 5, therefore place 5 under your second Prick, and fet 325 under 341, and substracting it from 341, there will re: main 16, which set over Head, then will the Sum appear thus:

54. The Art of Numbring

And in the middle Column

of your Square Rod against this 5, there stands

10; for this 10 you should
take a Rod that hath 10
at the Top, but being
there is no such, take
therefore one that hath a

Cypher, and place that between your Square-Rod and your Rod of 6, and change your Rod of 6 for one of 7, then

Thus must you al- | shall you have upways do, when the | on your Tabulat Number in the mid- | one Rod of 7, adle Column exceeds | nother of 0, and 10. | your Square Rod.

Then, looking upon your Sum, you shall find 1685 to your third Prick, look therefore upon your Rods for the nearest less Number, which you shall find to be 1404, against which stands 2 in the last Column, set 2 between the Lines under the third Prick, and 1404 under 1685, and, substracting it from 1685, there will remain 281, which place above, so will your Sum stand thus:

And because the Number, standing against in the middle Column of your Square Rod between 1404, and 2 was 4, set 4 under your last Prick, and take a Rod of 4, and put it between your Square Rod, and your Rod of 0; and because 28176 remains upon your Sum to the last

Prick, look upon your Rods for the nearest Number thereunto, and you shall find the very Number it self to stand against the Figure 4, set therefore 28176 below, and substract it from that above, and there will remain nothing, which denotes the Number 12418576, to be a square Number, and the Root thereof to be 3524, and the Work sinish'd will stand thus:

36 The Art of Numbring

This Sum, had it been wrought by the fecond Way of Division, which I shewed in Chapter 7, it would have stood as followeth:

Square 12418576 (3524 Root.

Caution.

Caution

If at any Time you look for the Remainder upon your Rods, and you cannot find it there, you must then place a Cypher between the Lines, and proceed to the next Figure, as by trying this other Example, which I have inserted for Practice, will appear.

Another Example added for Practice.

CHAP, X.

Concerning the Extraction of the Cube-Root.

HERE is somewhat more Difficulty in Extracting of the Cube, than of the Square Root. Wherefore (before I come to an Example) I will deliver the Manner of the Operation, together with fuch Cautions as are to be observed in the Performance thereof; all which immediately follow in this

GENERAL RULE.

Write down the Number whose Cube Root you are to extract, and under the first Figure towards the right Hand make a Prick or Point, and so under every third Figure towards the left Hand, till you come to the end of your Number. Under these Pricks draw two parallel Lines, (as you did in Extrasting the Square Root) between which Lines you are to place the Figures of your Root as you find

find them: — Then beginning at the Figure (or Figures) of the left Hand Prick, and going forward towards the right Hand, extract (by Help of the Rod for Extracting the Cube Root) their Root, or, if the true Number be not on the Plate, then the nearest less, and, placing this Root (which never exceeds one Figure) between the Lines, and under its Point, take its Cube from the uppermost Figure, which stands before (or leftwards of) the first Point, and note the Remainder above.

Secondly, Keep the Triple of this Root found, in the Head or Top of the Rods, and triple the Square of the same Root, and set this Triple on the Head of the Rods, and apply it leftwards of the Cubick Rod, and the reserved Rod (or Rods) rightwards, the Cubick Rod being in the midst between them; and out of the left Hand Rods, and the Cubick Rod together, pick or find out the Multiple, (or next less Number than the Figures preceding the second Point) which write apart in a Paper, and note its Quotum over its utmost right Hand Figure, and write the Square of that Quotum leftwards from the Quotum itself, even in that Order as you find them in your Cubick Rod; and N 2 under

under every several Figure of this Square, write their Multiples found rightwards, even such as the Figures themselves do shew; so that every Multiple may end under its Figure or Quotum: Then add together these Multiples cross-wise, and take their Sum from the Figures foregoing the second Point, and write the Remainder over them; but write the right Hand Quotum, before noted, under the second Point between the Lines, for the second Figure or Quotum of the Root: And so is performed the Operation of the second Point, which you must repeat through the several Points, even to the last.

But in the Practice by this Rule, you may fometimes be at a stand, wherefore to this GENERAL RULE (that there may be no Obstacle) I will add

these two CAUTIONS.

CAUTION I.

But in all Operations and Points it must be observed, That if no Multiple (no not the least of all) found in the lest Rods, and the Plate, can be substracted from the foregoing Remains; then a Cypher [0] must be put under that Point for

the Quotum, the Remains being untouched, and abiding as before.

CAUTION II.

And if the aforesaid Sum, to be taken away, cannot be taken from the Figures going before its Point, the smaller Multiples must be added, which the next upper Quotums in the Cubick Rod do shew in the Rods, whose Sum may be taken away therefrom.

Example of the Cubick Extraction.

Let 22022635627 be a Number given, whose Cube Root you desire: Set down your Number, and point it, (beginning at 7, the last Figure towards the right Hand, and so under every third Figure) and draw two parallel Lines under it, and it will stand in this Manner;

22022635627

Look in your Rod, for the Extracting the Cube Root, for the nearest Cube Root of the Figures of your given Number, standing before the first Point towards your left Hand, namely, for the nearest Cube Root of the Number

less than 22, which you shall find to be 2, which fet between the two Lines just under the first Point, and its Cube (which is 8) fet under the Line, and substract it from the Figures above the Line, namely, from 22, and there will remain 14, which place orderly above, then will your Work stand thus, and the Work of your first Point finished;

> 14 22022635627

Secondly, For the finding of the Root belonging to the fecond Prick, triple the Quotum, or Figure which is under the first Prick, (namely 2) and it is 6; find therefore a Rod which hath 6 at the Head thereof, and lay that Rod by the Side of your Cubick Rod towards the right Hand, then triple the Square of 2 (which is 4) and it makes 12, which, found among the Rods, place by the Side of the Cubick Rod towards the left Hand.

Then from the Rods which lie on the left Hand of the Cubick Rod, and the Cubick

Cubick Rod itself, find the nearest lesser Number than the Figures standing before the second Prick, namely, less than 14022, and in the ninth Place you shall find 11529, which write by itself, as in the Margin, and over 9, the last Figure towards the right Hand (drawing first a Line between) set its Quotum, and by it its Square 81, in the same Order as you find them stand

in your Cubick Rod.

Then write under 1, its Multiple, which is shewed rightwards against i in the Cubick Rod, and is the fingle Figure 6, and under 8 write the Multiple, that it shews rightward against 8 in the Cubick Rod, which is 48, and these three Multiples so written cross-wise below the Line, and added together (as in the Margin) do produce 16389, which, because they cannot be taken from the upper Figures standing before the fecond Point, namely, from 14022, the Number 9 (before taken) is to be rejected, as being too great, and instead of 819 (by the second Caution) the next higher Notes in the Plate are to be taken, which are 648, and the Mul-

64 The Art of Numbring

Multiples that these do shew, namely, the Octuple among the left Rods, which is 10112, and the Quadruple among the right Rods, which is 24, and the Sextuple

among the right Rods, which is 36, being added cross-wife 648 (as in the Margin) do pro-10112 duce 13952, which substract. ed from 14022, (the Figures standing before the second F3952 Prick) there remains 70 for the Remainder of the second Prick, and let there be taken for the Quotum of the fecond Prick, the right-most of the chosen Figures 648, which is 8, which place under the second Point between the Lines; so is the second Figure of your Root found, and your Work will stand thus;

Thirdly, Put the Triple of the precedent Quotums (viz. 28 between the Lines) which is 48, being taken out of the Rods, and put them on the right Side of the Cubick Rod, and get the Triple of the Square of the same 28, which may be found to be 2352, which

take out of the Rods, and place on the left Side of the CubickRod; and of the Multiples on the left Hand Rods, and the simple single Figures upon the Cubick Rod (the least of which being 235201)

2325 there is none so little that may be substracted from the Figures belonging to the third Point, namely, from 70635: Therefore (by the first Caution) the Remains abiding, or continuing as they are, you must put a Cypher under the third Point, for the third Quotum belonging to the third Point: And thus the Operation of the third Point is accomplished, and the Work will stand as tolloweth;

Fourthly, Set the Triple of the foregoing Quotums (viz. 280) which is 840 on the right Hand, and the Triple of the Square of the same 280, which is 225200 on the left Hand, 280 and the Cubick Rod between 280 them; then, out of the left-22400 most Multiples, choose that 560 which is next less than the Fi-78400. gures belonging to the fourth Point, namely, 70635627, 235200. which is this 70560027, which stands against 3 on the Tabulat; wherefore write this Number 70560027 upon Paper, in the Margin, with a Line over it, and fet over the Line the .

right-most Figure, and the Square of the said Quotum 3, which is 9, lettward thereof, and the

93 70560027 7560 70635727

Noncuple found in the right Hand Rods, which is 7560, write under 9; let these two Multiples be added as in the Margin, and the Sum will be 70635617, which substracted from the Figures foregoing the fourth Prick, and there will nothing remain; therefore let the right-most of the Figures of 93, viz. 3, be placed under the fourth and last Point, for the fourth and last Quotum of the Root, and so the whole and perfect Cubick Root of the given Number 22022635627, is 2803, and, since nothing remained, it is a perfect Cubick Number. The like is to be done in other Numbers, but I Thall forbear to give you any more Examples, there falling out in this all the Variety that at any Time may happen for the General Rule; and the two Cautions before premised are here made applicable to Practice; wherefore to this Treatise for the present I will put

FINIS.

Page 12, Line 25, for 3 Degr. read 31 d. 46 m.

ADVERTISEMENT.

Compleat System of Astronomy. In Two Volumes. Containing the Description and Use of the Sector; the Laws of Spheric Geometry; the Projestion of the Sphere Orthographically and Stereographically upon the Planes of the Meridian, Ecliptic and Horizon; the Doctrine of the Sphere; and the E; clipses of the Sun and Moon for thirty feven Years. Together with all the Precepts of Calculation. Also new Tables of the Motions of the Planets, fix'd Stars, and the first Satellite of Jupiter; of right and oblique Ascensions, and of logistical Logarithms. To the whole are prefix'd Astronomical Definitions for the Benefit of young Students. By Charles Leadbetter, Teacher of the Mathematicks, Printed for J. Wilcox, at the Green Dragon, in Little Britain.

Where may also be had,

The modern Navigator's compleat

Tutor, by J. Kelly, Mariner. Pr. 4s.
A new Treatise of the Construction and Use of the Sector, by Sam. Cun; revised by Ed. Stone. Pr. 3 s. 6 d.

that point of the Equator must necessarily by entightened thro The shadow which becomes it. To first the berson who makes use of the King Iral, Ela Cicle be drawn whon paper, equal in beameter by Aperture, which one judges proper to gue to the Bridge The Length of that Aperture is Equal to our Arch of the Meridian of 47 begree, to take in all of declinations of the Jun; & y Cicle drawn, which has of Aperture for a Viamoles, represent of Beliptic with its twelve Houses. What Rile, therefore, is divided into twelve equal parts, y bounds, thereof are joined two a two by parallel lines. which makes Spaces, narrower howards the Iropies, & Never bowards The Equinoscies, which lerve for hoslive Months, are divided afterwards into three Times to londays,

or into six Genes five, to suit, as near as possible, & position of the Cursor to the actual Tellination all their meaning use care fully transferred ony two Edges of the Bridge when this Instrument is intended for use, the tursor is put by day of the month, and the Suspension to the Latitude of the place: we turn, ofterwards, the Hat lide of the Bridge, wwards you Jun shis hays fall escarlly on the Edge of the Equator, except in the days of the Equinoxes, when the hun hirring round the brops Equator, as he does round y Celestial, can only east the Thodow of the upper Edga, on that opposite to it, the Hour of 12 of noon must alfo be excepted; breause the hen, shining then on the bross Meridian, thron the shadow thereof upon the opposit Edge, where, when yo glow of iz is marked one may know, however,

hat it is Mid-day, by the bial aving no radiant point at all

